

Inflation and the Theory of Cosmological Perturbations

A. Riotto^{†*}

[†] *INFN, Sezione di Padova, via Marzolo 8, I-35131, Padova, Italy.*

Lectures given at the:

Summer School on

Astroparticle Physics and Cosmology

Trieste, 17 June - 5 July 2002

Abstract

These lectures provide a pedagogical introduction to inflation and the theory of cosmological perturbations generated during inflation which are thought to be the origin of structure in the universe.

Keywords: Early Universe Physics and Particle Physics.

PACS numbers: 98.80.Cq

DFPD-TH/02/22

Contents

1	Introduction	1
2	Basics of the Big-Bang Model	1
2.1	Friedmann equations	2
2.2	Some conformalities	4
2.3	The early, radiation-dominated universe	5
2.4	The concept of particle horizon	7
3	The shortcomings of the Standard Big-Bang Theory	8
3.1	The Flatness Problem	8
3.2	The Entropy Problem	9
3.3	The horizon problem	9
4	The standard inflationary universe	13
4.1	Inflation and the horizon Problem	14
4.2	Inflation and the flatness problem	15
4.3	Inflation and the entropy problem	16
4.4	Inflation and the inflaton	16
4.5	Slow-roll conditions	17
4.6	The last stage of inflation and reheating	20
4.7	A brief survey of inflationary models	21
4.7.1	Large-field models	21
4.7.2	Small-field models	22
4.7.3	Hybrid models	22
5	Inflation and the cosmological perturbations	22
6	Quantum fluctuations of a generic massless scalar field during inflation	25
6.1	Quantum fluctuations of a generic massless scalar field during a de Sitter stage	25
6.2	Quantum fluctuations of a generic massive scalar field during a de Sitter stage	28
6.3	Quantum to classical transition	29
6.4	The power spectrum	30
6.5	Quantum fluctuations of a generic scalar field in a quasi de Sitter stage . . .	30
7	Quantum fluctuations during inflation	32
7.1	The metric fluctuations	34
7.2	Perturbed affine connections and Einstein's tensor	36
7.3	Perturbed stress energy-momentum tensor	39
7.4	Perturbed Klein-Gordon equation	40
7.5	The issue of gauge invariance	41
7.6	The comoving curvature perturbation	44
7.7	The curvature perturbation on spatial slices of uniform energy density	45
7.8	Scalar field perturbations in the spatially flat gauge	46
7.9	Adiabatic and isocurvature perturbations	47
7.10	The next steps	48

7.11	Computation of the curvature perturbation using the longitudinal gauge . .	49
7.12	Gauge-invariant computation of the curvature perturbation	52
7.13	Gravitational waves	57
7.14	The consistency relation	58
8	The post-inflationary evolution of the cosmological perturbations	59
8.1	From the inflationary seeds to the matter power spectrum	60
8.2	From inflation to large-angle CMB anisotropy	62
9	Conclusions	65
A	Evolution of the curvature perturbation on superhorizon scales	67
	References	68

1 Introduction

One of the basic ideas of modern cosmology is that there was an epoch early in the history of the universe when potential, or vacuum, energy dominated other forms of energy density such as matter or radiation. During the vacuum-dominated era the scale factor grew exponentially (or nearly exponentially) in time. In this phase, known as inflation, a small, smooth spatial region of size less than the Hubble radius at that time can grow so large as to easily encompass the comoving volume of the entire presently observable universe. If the early universe underwent this period of rapid expansion, then one can understand why the observed universe is so homogeneous and isotropic to high accuracy. All these virtues of inflation were noted when it was first proposed by Guth in 1981 [26]. A more dramatic consequence of the inflationary paradigm was noticed soon after [29, 63, 27]. Starting with a universe which is absolutely homogeneous and isotropic at the classical level, the inflationary expansion of the universe will ‘freeze in’ the vacuum fluctuation of the inflaton field so that it becomes an essentially classical quantity. On each comoving scale, this happens soon after horizon exit. Associated with this vacuum fluctuation is a primordial energy density perturbation, which survives after inflation and may be the origin of all structure in the universe. In particular, it may be responsible for the observed cosmic microwave background (CMB) anisotropy and for the large-scale distribution of galaxies and dark matter. Inflation also generates primordial gravitational waves as a vacuum fluctuation, which may contribute to the low multipoles of the CMB anisotropy. Therefore, a prediction of inflation is that all of the structure we see in the universe is a result of quantum-mechanical fluctuations during the inflationary epoch.

The goal of these lectures is to provide a pedagogical introduction to the inflationary paradigm and to the theory of cosmological perturbations generated during inflation.

The lectures are organized as follows. In section 2 we give a brief review of the Big-Bang theory. In section 3 we describe the shortcomings of the Big-Bang theory leading to the idea of inflation which is addressed in section 4. In section 5 we present the idea of quantum fluctuations which are studied in sections 6 and 7. Section 8 presents some details about the post-inflationary evolution of the cosmological perturbations. Finally, section 8 contains some conclusions.

Since these lectures were devoted to a school, we will not provide an exhaustive list of references to original material, but refer to several basic papers (including several review papers) where students can find the references to the original material. The list of references include Refs. [41, 36, 7, 64, 53, 38, 40, 48, 39].

Finally, we warn the reader that references are listed in alphabetical order.

2 Basics of the Big-Bang Model

The standard cosmology is based upon the maximally spatially symmetric Friedmann-Robertson-Walker (FRW) line element

$$ds^2 = -dt^2 + a(t)^2 \left[\frac{dr^2}{1 - kr^2} + r^2(d\theta^2 + \sin^2 \theta d\phi^2) \right]; \quad (1)$$

where $a(t)$ is the cosmic-scale factor, $R_{\text{curv}} \equiv a(t)|k|^{-1/2}$ is the curvature radius, and $k = -1, 0, 1$ is the curvature signature. All three models are without boundary: the positively curved model is finite and “curves” back on itself; the negatively curved and flat models are infinite in extent. The Robertson-Walker metric embodies the observed isotropy and homogeneity of the universe. It is interesting to note that this form of the line element was originally introduced for sake of mathematical simplicity; we now know that it is well justified at early times or today on large scales ($\gg 10$ Mpc), at least within our visible patch.

The coordinates, r , θ , and ϕ , are referred to as *comoving* coordinates: A particle at rest in these coordinates remains at rest, *i.e.*, constant r , θ , and ϕ . A freely moving particle eventually comes to rest these coordinates, as its momentum is red shifted by the expansion, $p \propto a^{-1}$. Motion with respect to the comoving coordinates (or cosmic rest frame) is referred to as peculiar velocity; unless “supported” by the inhomogeneous distribution of matter peculiar velocities decay away as a^{-1} . Thus the measurement of peculiar velocities, which is not easy as it requires independent measures of both the distance and velocity of an object, can be used to probe the distribution of mass in the universe.

Physical separations between freely moving particles scale as $a(t)$; or said another way the physical separation between two points is simply $a(t)$ times the coordinate separation. The momenta of freely propagating particles decrease, or “red shift,” as $a(t)^{-1}$, and thus the wavelength of a photon stretches as $a(t)$, which is the origin of the cosmological red shift. The red shift suffered by a photon emitted from a distant galaxy $1 + z = a_0/a(t)$; that is, a galaxy whose light is red shifted by $1 + z$, emitted that light when the universe was a factor of $(1 + z)^{-1}$ smaller. Thus, when the light from the most distant quasar yet seen ($z = 4.9$) was emitted the universe was a factor of almost six smaller; when CMB photons last scattered the universe was about 1100 times smaller.

2.1 Friedmann equations

The evolution of the scale factor $a(t)$ is governed by Einstein equations

$$R_{\mu\nu} - \frac{1}{2} R g_{\mu\nu} \equiv G_{\mu\nu} = 8\pi G T_{\mu\nu} \quad (2)$$

where $R_{\mu\nu}$ ($\mu, \nu = 0, \dots, 3$) is the Riemann tensor and R is the Ricci scalar constructed via the metric (1) [36] and $T_{\mu\nu}$ is the energy-momentum tensor. Under the hypothesis of homogeneity and isotropy, we can always write the energy-momentum tensor under the form $T_{\mu\nu} = \text{diag}(\rho, p, p, p)$ where ρ is the energy density of the system and p its pressure. They are functions of time.

The evolution of the cosmic-scale factor is governed by the Friedmann equation

$$H^2 \equiv \left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G \rho}{3} - \frac{k}{a^2}; \quad (3)$$

where ρ is the total energy density of the universe, matter, radiation, vacuum energy, and so on.

Differentiating wrt to time both members of Eq. (177) and using the the mass conservation equation

$$\dot{\rho} + 3H(\rho + p) = 0, \quad (4)$$

we find the equation for the acceleration of the scale-factor

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3}(\rho + 3p). \quad (5)$$

Combining Eqs. (3) and (5) we find

$$\dot{H} = -4\pi G(\rho + p). \quad (6)$$

The evolution of the energy density of the universe is governed by

$$d(\rho a^3) = -pd(a^3); \quad (7)$$

which is the First Law of Thermodynamics for a fluid in the expanding universe. (In the case that the stress energy of the universe is comprised of several, noninteracting components, this relation applies to each separately; *e.g.*, to the matter and radiation separately today.) For $p = \rho/3$, ultra-relativistic matter, $\rho \propto a^{-4}$ and $a \sim t^{\frac{1}{2}}$; for $p = 0$, very nonrelativistic matter, $\rho \propto a^{-3}$ and $a \sim t^{\frac{2}{3}}$; and for $p = -\rho$, vacuum energy, $\rho = \text{const.}$ If the rhs of the Friedmann equation is dominated by a fluid with equation of state $p = \gamma\rho$, it follows that $\rho \propto a^{-3(1+\gamma)}$ and $a \propto t^{2/3(1+\gamma)}$.

We can use the Friedmann equation to relate the curvature of the universe to the energy density and expansion rate:

$$\Omega - 1 = \frac{k}{a^2 H^2}; \quad \Omega = \frac{\rho}{\rho_{\text{crit}}}; \quad (8)$$

and the critical density today $\rho_{\text{crit}} = 3H^2/8\pi G = 1.88h^2 \text{ g cm}^{-3} \simeq 1.05 \times 10^4 \text{ eV cm}^{-3}$. There is a one to one correspondence between Ω and the spatial curvature of the universe: positively curved, $\Omega_0 > 1$; negatively curved, $\Omega_0 < 1$; and flat ($\Omega_0 = 1$). Further, the “fate of the universe” is determined by the curvature: model universes with $k \leq 0$ expand forever, while those with $k > 0$ necessarily recollapse. The curvature radius of the universe is related to the Hubble radius and Ω by

$$R_{\text{curv}} = \frac{H^{-1}}{|\Omega - 1|^{1/2}}. \quad (9)$$

In physical terms, the curvature radius sets the scale for the size of spatial separations where the effects of curved space become “pronounced.” And in the case of the positively curved model it is just the radius of the 3-sphere.

The energy content of the universe consists of matter and radiation (today, photons and neutrinos). Since the photon temperature is accurately known, $T_0 = 2.73 \pm 0.01 \text{ K}$, the fraction of critical density contributed by radiation is also accurately known: $\Omega_R h^2 = 4.2 \times 10^{-5}$, where $h = 0.72 \pm 0.07$ is the present Hubble rate in units of $100 \text{ km sec}^{-1} \text{ Mpc}^{-1}$ [25]. The remaining content of the universe is another matter. Rapid progress has been made recently toward the measurement of cosmological parameters [6]. Over the past three years the basic features of our universe have been determined. The universe is spatially flat; accelerating; comprised of one third of dark matter and two third a new form of dark energy. The measurements of the cosmic microwave background anisotropies at different angular scales performed by Boomerang, Maxima, DASI, CBI and VSA have recently significantly

increase the case for accelerated expansion in the early universe (the inflationary paradigm) and at the current epoch (dark energy dominance), especially when combined with data on high redshift supernovae (SN1) and large scale structure (LSS) [6]. A recent analysis [14] shows that the CMB+LSS+SN1 data give

$$\Omega_0 = 1.00^{+0.07}_{-0.03},$$

meaning tha the present universe is spatially flat (or at least very close to being flat). Restricting to $\Omega_0 = 1$, the dark matter density is given by [14]

$$\Omega_{\text{DM}} h^2 = 0.12^{+0.01}_{-0.01},$$

and a baryon density

$$\Omega_B h^2 = 0.022^{+0.003}_{-0.002},$$

while the Big Bang nucleosynthesis estimate is $\Omega_B h^2 = 0.019 \pm 0.002$. Substantial dark (unclustered) energy is inferred,

$$\Omega_Q \approx 0.68 \pm 0.05,$$

compatible with the independent SN1 estimates! What is most relevant for us, this universe is apparently born from a burst of rapid expansion, inflation, during which quantum noise was stretched to astrophysical size seeding cosmic structure. This is exactly the phenomena we want to address in these lectures.

2.2 Some conformalities

Before launching ourselves into the description of the early universe, we would like to introduce the concept of conformal time which will be useful in the next sections. The conformal time τ is defined through the following relation

$$d\tau = \frac{dt}{a}. \quad (10)$$

The metric (1) then becomes

$$ds^2 = -a^2(\tau) \left[d\tau^2 - \frac{dr^2}{1 - kr^2} - r^2(d\theta^2 + \sin^2 \theta d\phi^2) \right]. \quad (11)$$

The reason why τ is called conformal is manifest from Eq. (11): the corresponding FRW line element is conformal to the Minkowski line element describing a static four dimensional hypersurface.

Any function $f(t)$ satisfies the rule

$$\dot{f}(t) = \frac{f'(\tau)}{a(\tau)}, \quad (12)$$

$$\ddot{f}(t) = \frac{f''(\tau)}{a^2(\tau)} - \mathcal{H} \frac{f'(\tau)}{a^2(\tau)}, \quad (13)$$

where a prime now indicates differentiation wrt to the conformal time τ and

$$\mathcal{H} = \frac{a'}{a}. \quad (14)$$

In particular we can set the following rules

$$\begin{aligned} H &= \frac{\dot{a}}{a} = \frac{a'}{a^2} = \frac{\mathcal{H}}{a}, \\ \ddot{a} &= \frac{a''}{a^2} - \frac{\mathcal{H}^2}{a}, \\ \dot{H} &= \frac{\mathcal{H}'}{a^2} - \frac{\mathcal{H}^2}{a^2}, \\ H^2 &= \frac{8\pi G\rho}{3} - \frac{k}{a^2} \implies \mathcal{H}^2 = \frac{8\pi G\rho a^2}{3} - k \\ \dot{H} &= -4\pi G(\rho + p) \implies \mathcal{H}' = -\frac{4\pi G}{3}(\rho + 3p)a^2, \\ \dot{\rho} + 3H(\rho + p) &= 0 \implies \rho' + 3\mathcal{H}(\rho + p) = 0 \end{aligned}$$

Finally, if the scale factor $a(t)$ scales like $a \sim t^n$, solving the relation (10) we find

$$a \sim t^n \implies a(\tau) \sim \tau^{\frac{n}{1-n}}. \quad (15)$$

2.3 The early, radiation-dominated universe

In any case, at present, matter outweighs radiation by a wide margin. However, since the energy density in matter decreases as a^{-3} , and that in radiation as a^{-4} (the extra factor due to the red shifting of the energy of relativistic particles), at early times the universe was radiation dominated—indeed the calculations of primordial nucleosynthesis provide excellent evidence for this. Denoting the epoch of matter-radiation equality by subscript ‘EQ,’ and using $T_0 = 2.73$ K, it follows that

$$a_{\text{EQ}} = 4.18 \times 10^{-5} (\Omega_0 h^2)^{-1}; \quad T_{\text{EQ}} = 5.62 (\Omega_0 h^2) \text{ eV}; \quad (16)$$

$$t_{\text{EQ}} = 4.17 \times 10^{10} (\Omega_0 h^2)^{-2} \text{ sec}. \quad (17)$$

At early times the expansion rate and age of the universe were determined by the temperature of the universe and the number of relativistic degrees of freedom:

$$\rho_{\text{rad}} = g_*(T) \frac{\pi^2 T^4}{30}; \quad H \simeq 1.67 g_*^{1/2} T^2 / m_{\text{Pl}}; \quad (18)$$

$$\implies a \propto t^{1/2}; \quad t \simeq 2.42 \times 10^{-6} g_*^{-1/2} (T / \text{GeV})^{-2} \text{ sec}; \quad (19)$$

where $g_*(T)$ counts the number of ultra-relativistic degrees of freedom ($\approx$ the sum of the internal degrees of freedom of particle species much less massive than the temperature) and $m_{\text{Pl}} \equiv G^{-1/2} = 1.22 \times 10^{19} \text{ GeV}$ is the Planck mass. For example, at the epoch of nucleosynthesis, $g_* = 10.75$ assuming three, light ($\ll \text{ MeV}$) neutrino species; taking into account all the species in the standard model, $g_* = 106.75$ at temperatures much greater than 300 GeV .

A quantity of importance related to g_* is the entropy density in relativistic particles,

$$s = \frac{\rho + p}{T} = \frac{2\pi^2}{45} g_* T^3,$$

and the entropy per comoving volume,

$$S \propto a^3 s \propto g_* a^3 T^3.$$

By a wide margin most of the entropy in the universe exists in the radiation bath. The entropy density is proportional to the number density of relativistic particles. At present, the relativistic particle species are the photons and neutrinos, and the entropy density is a factor of 7.04 times the photon-number density: $n_\gamma = 413 \text{ cm}^{-3}$ and $s = 2905 \text{ cm}^{-3}$.

In thermal equilibrium—which provides a good description of most of the history of the universe—the entropy per comoving volume S remains constant. This fact is very useful. First, it implies that the temperature and scale factor are related by

$$T \propto g_*^{-1/3} a^{-1}, \tag{20}$$

which for $g_* = \text{const}$ leads to the familiar $T \propto a^{-1}$.

Second, it provides a way of quantifying the net baryon number (or any other particle number) per comoving volume:

$$N_B \equiv R^3 n_B = \frac{n_B}{s} \simeq (4 - 7) \times 10^{-11}. \tag{21}$$

The baryon number of the universe tells us two things: (1) the entropy per particle in the universe is extremely high, about 10^{10} or so compared to about 10^{-2} in the sun and a few in the core of a newly formed neutron star. (2) The asymmetry between matter and antimatter is very small, about 10^{-10} , since at early times quarks and antiquarks were roughly as abundant as photons. One of the great successes of particle cosmology is baryogenesis, the idea that B , C , and CP violating interactions occurring out-of-equilibrium early on allow the universe to develop a net baryon number of this magnitude [59, 60].

Finally, the constancy of the entropy per comoving volume allows us to characterize the size of comoving volume corresponding to our present Hubble volume in a very physical way: by the entropy it contains,

$$S_U = \frac{4\pi}{3} H_0^{-3} s \simeq 10^{90}. \tag{22}$$

The standard cosmology is tested back to times as early as about 0.01 sec; it is only natural to ask how far back one can sensibly extrapolate. Since the fundamental particles of Nature are point-like quarks and leptons whose interactions are perturbatively weak at energies much greater than 1 GeV , one can imagine extrapolating as far back as the epoch

where general relativity becomes suspect, i.e., where quantum gravitational effects are likely to be important: the Planck epoch, $t \sim 10^{-43}$ sec and $T \sim 10^{19}$ GeV. Of course, at present, our firm understanding of the elementary particles and their interactions only extends to energies of the order of 100 GeV, which corresponds to a time of the order of 10^{-11} sec or so. We can be relatively certain that at a temperature of 100 MeV–200 MeV ($t \sim 10^{-5}$ sec) there was a transition (likely a second-order phase transition) from quark/gluon plasma to very hot hadronic matter, and that some kind of phase transition associated with the symmetry breakdown of the electroweak theory took place at a temperature of the order of 300 GeV ($t \sim 10^{-11}$ sec).

2.4 The concept of particle horizon

In spite of the fact that the universe was vanishingly small at early times, the rapid expansion precluded causal contact from being established throughout. Photons travel on null paths characterized by $dr = dt/a(t)$; the physical distance that a photon could have traveled since the bang until time t , the distance to the particle horizon, is

$$\begin{aligned} R_H(t) &= a(t) \int_0^t \frac{dt'}{a(t')} \\ &= \frac{t}{(1-n)} = n \frac{H^{-1}}{(1-n)} \sim H^{-1} \quad \text{for } a(t) \propto t^n, \quad n < 1. \end{aligned} \quad (23)$$

Using the conformal time, the particle horizon becomes

$$R_H(t) = a(\tau) \int_{\tau_0}^{\tau} d\tau, \quad (24)$$

where τ_0 indicates the conformal time corresponding to $t = 0$. Note, in the standard cosmology the distance to the horizon is finite, and up to numerical factors, equal to the age of the universe or the Hubble radius, H^{-1} . For this reason, we will use horizon and Hubble radius interchangeably.¹

Note also that a physical length scale λ is within the horizon if $\lambda < R_H \sim H^{-1}$. Since we can identify the length scale λ with its wavenumber k , $\lambda = 2\pi a/k$, we will have the following rule

$\begin{aligned} \frac{k}{aH} &\ll 1 \implies \text{SCALE } \lambda \text{ OUTSIDE THE HORIZON} \\ \frac{k}{aH} &\gg 1 \implies \text{SCALE } \lambda \text{ WITHIN THE HORIZON} \end{aligned}$

¹As we shall see, in inflationary models the horizon and Hubble radius are not roughly equal as the horizon distance grows exponentially relative to the Hubble radius; in fact, at the end of inflation they differ by e^N , where N is the number of e-folds of inflation. However, we will slip and use “horizon” and “Hubble radius” interchangeably, though we will always mean Hubble radius.

An important quantity is the entropy within a horizon volume: $S_{\text{HOR}} \sim H^{-3}T^3$; during the radiation-dominated epoch $H \sim T^2/m_{\text{Pl}}$, so that

$$S_{\text{HOR}} \sim \left(\frac{m_{\text{Pl}}}{T}\right)^3; \quad (25)$$

from this we will shortly conclude that at early times the comoving volume that encompasses all that we can see today (characterized by an entropy of about 10^{90}) was comprised of a very large number of causally disconnected regions.

3 The shortcomings of the Standard Big-Bang Theory

By now the shortcomings of the standard cosmology are well appreciated: the horizon or large-scale smoothness problem; the small-scale inhomogeneity problem (origin of density perturbations); and the flatness or oldness problem. We will only briefly review them here. They do not indicate any logical inconsistencies of the standard cosmology; rather, that very special initial data seem to be required for evolution to a universe that is qualitatively similar to ours today. Nor is inflation the first attempt to address these shortcomings: over the past two decades cosmologists have pondered this question and proposed alternative solutions. Inflation is a solution based upon well-defined, albeit speculative, early universe microphysics describing the post-Planck epoch.

3.1 The Flatness Problem

Let us make a tremendous extrapolation and assume that Einstein equations are valid until the Planck era, when the temperature of the universe is $T_{\text{Pl}} \sim m_{\text{Pl}} \sim 10^{19}$ GeV. From Eq. (8), we read that if the universe is perfectly flat, then $(\Omega = 1)$ at all times. On the other hand, if there is even a small curvature term, the time dependence of $(\Omega - 1)$ is quite different.

During a radiation-dominated period, we have that $H^2 \propto \rho_R \propto a^{-4}$ and

$$\Omega - 1 \propto \frac{1}{a^2 a^{-4}} \propto a^2. \quad (26)$$

During Matter Domination, $\rho_M \propto a^{-3}$ and

$$\Omega - 1 \propto \frac{1}{a^2 a^{-3}} \propto a. \quad (27)$$

In both cases $(\Omega - 1)$ decreases going backwards with time. Since we know that today $(\Omega_0 - 1)$ is of order unity at present, we can deduce its value at t_{Pl} (the time at which the temperature of the universe is $T_{\text{Pl}} \sim 10^{19}$ GeV)

$$\frac{|\Omega - 1|_{T=T_{\text{Pl}}}}{|\Omega - 1|_{T=T_0}} \approx \left(\frac{a_{\text{Pl}}^2}{a_0^2}\right) \approx \left(\frac{T_0^2}{T_{\text{Pl}}^2}\right) \approx \mathcal{O}(10^{-64}). \quad (28)$$

where 0 stands for the present epoch, and $T_0 \sim 10^{-13}$ GeV is the present-day temperature of the CMB radiation. If we are not so brave and go back simply to the epoch of nucleosynthesis when light elements abundances were formed, at $T_N \sim 1$ MeV, we get

$$\frac{|\Omega - 1|_{T=T_N}}{|\Omega - 1|_{T=T_0}} \approx \left(\frac{a_N^2}{a_0^2}\right) \approx \left(\frac{T_0^2}{T_N^2}\right) \approx \mathcal{O}(10^{-16}). \quad (29)$$

In order to get the correct value of $(\Omega_0 - 1) \sim 1$ at present, the value of $(\Omega - 1)$ at early times have to be fine-tuned to values amazingly close to zero, but without being exactly zero. This is the reason why the flatness problem is also dubbed the ‘fine-tuning problem’.

3.2 The Entropy Problem

Let us now see how the hypothesis of adiabatic expansion of the universe is connected with the flatness problem. From the Friedman equation (177) we know that during a radiation-dominated period

$$H^2 \simeq \rho_R \simeq \frac{T^4}{m_{\text{Pl}}^2}, \quad (30)$$

from which we deduce

$$\Omega - 1 = \frac{km_{\text{Pl}}^2}{a^4 T^4} = \frac{km_{\text{Pl}}^2}{S^{\frac{2}{3}} T^2}. \quad (31)$$

Under the hypothesis of adiabaticity, S is constant over the evolution of the universe and therefore

$$|\Omega - 1|_{t=t_{\text{Pl}}} = \frac{m_{\text{Pl}}^2}{T_{\text{Pl}}^2} \frac{1}{S_U^{2/3}} = \frac{1}{S_U^{2/3}} \approx 10^{-60}. \quad (32)$$

We have discovered that $(\Omega - 1)$ is so close to zero at early epochs because the total entropy of our universe is so incredibly large. The flatness problem is therefore a problem of understanding why the (classical) initial conditions corresponded to a universe that was so close to spatial flatness. In a sense, the problem is one of fine-tuning and although such a balance is possible in principle, one nevertheless feels that it is unlikely. On the other hand, the flatness problem arises because the entropy in a comoving volume is conserved. It is possible, therefore, that the problem could be resolved if the cosmic expansion was non-adiabatic for some finite time interval during the early history of the universe.

3.3 The horizon problem

According to the standard cosmology, photons decoupled from the rest of the components (electrons and baryons) at a temperature of the order of 0.3 eV. This corresponds to the so-called surface of ‘last-scattering’ at a red shift of about 1100 and an age of about $180,000 (\Omega_0 h^2)^{-1/2}$ yrs. From the epoch of last-scattering onwards, photons free-stream and reach us basically untouched. Detecting primordial photons is therefore equivalent to take a picture of the universe when the latter was about 300,000 yrs old. The spectrum of the

cosmic background radiation (CBR) is consistent that of a black body at temperature 2.73 K over more than three decades in wavelength; see Fig. 1.

The most accurate measurement of the temperature and spectrum is that by the FIRAS instrument on the COBE satellite which determined its temperature to be 2.726 ± 0.01 K [49]. The length corresponding to our present Hubble radius (which is approximately the radius of our observable universe) at the time of last-scattering was

$$\lambda_H(t_{LS}) = R_H(t_0) \left(\frac{a_{LS}}{a_0} \right) = R_H(t_0) \left(\frac{T_0}{T_{LS}} \right).$$

On the other hand, during the matter-dominated period, the Hubble length has decreased with a different law

$$H^2 \propto \rho_M \propto a^{-3} \propto T^3.$$

At last-scattering

$$H_{LS}^{-1} = R_H(t_0) \left(\frac{T_{LS}}{T_0} \right)^{-3/2} \ll R_H(t_0).$$

The length corresponding to our present Hubble radius was much larger than the horizon at that time. This can be shown comparing the volumes corresponding to these two scales

$$\frac{\lambda_H^3(T_{LS})}{H_{LS}^{-3}} = \left(\frac{T_0}{T_{LS}} \right)^{-\frac{3}{2}} \approx 10^6. \quad (33)$$

There were $\sim 10^6$ casually disconnected regions within the volume that now corresponds to our horizon! It is difficult to come up with a process other than an early hot and dense phase in the history of the universe that would lead to a precise black body [55] for a bath of photons which were causally disconnected the last time they interacted with the surrounding plasma.

The horizon problem is well represented by Fig. 2 where the green line indicates the horizon scale and the red line any generic physical length scale λ . Suppose, indeed that λ indicates the distance between two photons we detect today. From Eq. (33) we discover that at the time of emission (last-scattering) the two photons could not talk to each other, the red line is above the green line. There is another aspect of the horizon problem which is related to the problem of initial conditions for the cosmological perturbations. We have every indication that the universe at early times, say $t \ll 300,000$ yrs, was very homogeneous; however, today inhomogeneity (or structure) is ubiquitous: stars ($\delta\rho/\rho \sim 10^{30}$), galaxies ($\delta\rho/\rho \sim 10^5$), clusters of galaxies ($\delta\rho/\rho \sim 10 - 10^3$), superclusters, or “clusters of clusters” ($\delta\rho/\rho \sim 1$), voids ($\delta\rho/\rho \sim -1$), great walls, and so on. For some twenty-five years the standard cosmology has provided a general framework for understanding this picture. Once the universe becomes matter dominated (around 1000 yrs after the bang) primeval density inhomogeneities ($\delta\rho/\rho \sim 10^{-5}$) are amplified by gravity and grow into the structure we see today [54]. The existence of density inhomogeneities has another important consequence: fluctuations in the temperature of the CMB radiation of a similar amplitude. The temperature difference measured between two points separated by a large angle ($\gtrsim 1^\circ$) arises due to a very simple physical effect: the difference in the gravitational potential between the two points on the last-scattering surface, which in turn is related to the density perturbation, determines the temperature anisotropy

on the angular scale subtended by that length scale,

$$\left(\frac{\delta T}{T}\right)_\theta \approx \left(\frac{\delta \rho}{\rho}\right)_\lambda, \quad (34)$$

where the scale $\lambda \sim 100h^{-1} \text{ Mpc}(\theta/\text{deg})$ subtends an angle θ on the last-scattering surface. This is known as the Sachs-Wolfe effect [61]. The CMB experiments looking for the tiny anisotropies are of three kinds: satellite experiments, balloon experiments, and ground based experiments. The technical and economical advantages of ground based experiments are evident, but their main problem is atmospheric fluctuations. The problem can be limited choosing a very high and cold site, or working on small scales (as the Dasi experiment [18]). Balloon based experiments limit the atmospheric problems, but have to face the following problems: they must be limited in weight, they can not be manipulated during the flight, they have a rather short duration (and they have to be recovered intact). Maxima [51], and Boomerang [15] are experiments of this kind.

At present, there is a satellite experiment – MAP (Microwave Anisotropy Probe) sponsored by NASA mission, which is taking data [50]. Finally, a satellite mission PLANCK is planned by ESA to be launched in 2007 [56]. The temperature anisotropy is commonly expanded in spherical harmonics

$$\frac{\Delta T}{T}(x_0, \tau_0, \mathbf{n}) = \sum_{\ell m} a_{\ell, m}(x_0) Y_{\ell m}(\mathbf{n}), \quad (35)$$

where x_0 and τ_0 are our position and the preset time, respectively, $\mathbf{n}$ is the direction of observation, ℓ 's are the different multipoles and²

$$\langle a_{\ell m} a_{\ell' m'}^* \rangle = \delta_{\ell, \ell'} \delta_{m, m'} C_\ell, \quad (36)$$

where the deltas are due to the fact that the process that created the anisotropy is statistically isotropic. The C_ℓ are the so-called CMB power spectrum. For homogeneity and isotropy, the C_ℓ 's are neither a function of x_0 , nor of m . The two-point-correlation function is related to the C_ℓ 's in the following way

$$\begin{aligned} \left\langle \frac{\delta T(\mathbf{n})}{T} \frac{\delta T(\mathbf{n}')}{T} \right\rangle &= \sum_{\ell \ell' m m'} \langle a_{\ell m} a_{\ell' m'}^* \rangle Y_{\ell m}(\mathbf{n}) Y_{\ell' m'}^*(\mathbf{n}') \\ &= \sum_{\ell} C_\ell \sum_m Y_{\ell m}(\mathbf{n}) Y_{\ell m}^*(\mathbf{n}') = \frac{1}{4\pi} \sum_{\ell} (2\ell + 1) C_\ell P_\ell(\mu = \mathbf{n} \cdot \mathbf{n}') \end{aligned} \quad (37)$$

where we have used the addition theorem for the spherical harmonics, and P_ℓ is the Legendre polynomial of order ℓ . In expression (37) the expectation value is an ensemble average. It can be regarded as an average over the possible observer positions, but not in general as an average over the single sky we observe, because of the cosmic variance³.

²An alternative definition is $C_\ell = \langle |a_{\ell m}|^2 \rangle = \frac{1}{2\ell+1} \sum_{m=-\ell}^{\ell} |a_{\ell m}|^2$.

³The usual hypothesis is that we observe a typical realization of the ensemble. This means that we expect the difference between the observed values $|a_{\ell m}|^2$ and the ensemble averages C_ℓ to be of the order of the mean-square deviation of $|a_{\ell m}|^2$ from C_ℓ . The latter is called cosmic variance and, because we are dealing with a gaussian distribution, it is equal to $2C_\ell$ for each multipole ℓ . For a single ℓ , averaging over the $(2\ell+1)$ values of m reduces the cosmic variance by a factor $(2\ell+1)$, but it remains a serious limitation for low multipoles.

Let us now consider the last-scattering surface. In comoving coordinates the latter is ‘far’ from us a distance equal to

$$\int_{t_{\text{LS}}}^{t_0} \frac{dt}{a} = \int_{\tau_{\text{LS}}}^{\tau_0} d\tau = (\tau_0 - \tau_{\text{LS}}). \quad (38)$$

A given comoving scale λ is therefore projected on the last-scattering surface sky on an angular scale

$$\theta \simeq \frac{\lambda}{(\tau_0 - \tau_{\text{LS}})}, \quad (39)$$

where we have neglected tiny curvature effects. Consider now that the scale λ is of the order of the comoving sound horizon at the time of last-scattering, $\lambda \sim c_s \tau_{\text{LS}}$, where $c_s \simeq 1/\sqrt{3}$ is the sound velocity at which photons propagate in the plasma at the last-scattering. This corresponds to an angle

$$\theta \simeq c_s \frac{\tau_{\text{LS}}}{(\tau_0 - \tau_{\text{LS}})} \simeq c_s \frac{\tau_{\text{LS}}}{\tau_0}, \quad (40)$$

where the last passage has been performed knowing that $\tau_0 \gg \tau_{\text{LS}}$. Since the universe is matter-dominated from the time of last-scattering onwards, the scale factor has the following behaviour: $a \sim T^{-1} \sim t^{2/3} \sim \tau^2$, where we have made use of the relation (15). The angle θ_{HOR} subtended by the sound horizon on the last-scattering surface then becomes

$$\theta_{\text{HOR}} \simeq c_s \left(\frac{T_0}{T_{\text{LS}}} \right)^{1/2} \sim 1^\circ, \quad (41)$$

where we have used $T_{\text{LS}} \simeq 0.3$ eV and $T_0 \sim 10^{-13}$ GeV. This corresponds to a multipole ℓ_{HOR}

$$\ell_{\text{HOR}} = \frac{\pi}{\theta_{\text{HOR}}} \simeq 200. \quad (42)$$

From these estimates we conclude that two photons which on the last-scattering surface were separated by an angle larger than θ_{HOR} , corresponding to multipoles smaller than $\ell_{\text{HOR}} \sim 200$ were not in causal contact. On the other hand, from Fig. (3) it is clear that small anisotropies, of the *same* order of magnitude $\delta T/T \sim 10^{-5}$ are present at $\ell \ll 200$. We conclude that one of the striking features of the CMB fluctuations is that they appear to be noncausal. Photons at the last-scattering surface which were causally disconnected have the same small anisotropies! The existence of particle horizons in the standard cosmology precludes explaining the smoothness as a result of microphysical events: the horizon at decoupling, the last time one could imagine temperature fluctuations being smoothed by particle interactions, corresponds to an angular scale on the sky of about 1° , which precludes temperature variations on larger scales from being erased.

To account for the small-scale lumpiness of the universe today, density perturbations with horizon-crossing amplitudes of 10^{-5} on scales of 1 Mpc to 10^4 Mpc or so are required. As can be seen in Fig. 2, in the standard cosmology the physical size of a perturbation, which grows as the scale factor, begins larger than the horizon and relatively late in the history of the universe crosses inside the horizon. This precludes a causal microphysical explanation for the origin of the required density perturbations.

From the considerations made so far, it appears that solving the shortcomings of the standard Big Bang theory requires two basic modifications of the assumptions made so far:

- The universe has to go through a non-adiabatic period. This is necessary to solve the entropy and the flatness problem. A non-adiabatic phase may give rise to the large entropy S_U we observe today.
- The universe has to go through a primordial period during which the physical scales λ evolve faster than the horizon scale H^{-1} .

The second condition is obvious from Fig. 4. If there is period during which physical length scales grow faster than H^{-1} , length scales λ which are within the horizon today, $\lambda < H^{-1}$ (such as the distance between two detected photons) and were outside the horizon for some period, $\lambda > H^{-1}$ (for instance at the time of last-scattering when the two photons were emitted), had a chance to be within the horizon at some primordial epoch, $\lambda < H^{-1}$ again. If this happens, the homogeneity and the isotropy of the CMB can be easily explained: photons that we receive today and were emitted from the last-scattering surface from causally disconnected regions have the same temperature because they had a chance to ‘talk’ to each other at some primordial stage of the evolution of the universe.

The second condition can be easily expressed as a condition on the scale factor a . Since a given scale λ scales like $\lambda \sim a$ and $H^{-1} = a/\dot{a}$, we need to impose that there is a period during which

$$\left(\frac{\lambda}{H^{-1}} \right)' = \ddot{a} > 0.$$

We can therefore introduced the following rigorous definition: an inflationary stage [26] is a period of the universe during which the latter accelerates

$$\text{INFLATION} \iff \ddot{a} > 0.$$

Comment: Let us stress that during such a accelerating phase the universe expands *adiabatically*. This means that during inflation one can exploit the usual FRW equations (3) and (5). It must be clear therefore that the non-adiabaticity condition is satisfied not during inflation, but during the phase transition between the end of inflation and the beginning of the radiation-dominated phase. At this transition phase a large entropy is generated under the form of relativistic degrees of freedom: the Big Bang has taken place.

4 The standard inflationary universe

From the previous section we have learned that an accelerating stage during the primordial phases of the evolution of the universe might be able to solve the horizon problem. From Eq. (5) we learn that

$$\ddot{a} > 0 \iff (\rho + 3p) < 0.$$

An accelerating period is obtainable only if the overall pressure p of the universe is negative: $p < -\rho/3$. Neither a radiation-dominated phase nor a matter-dominated phase (for which

$p = \rho/3$ and $p = 0$, respectively) satisfy such a condition. Let us postpone for the time being the problem of finding a ‘candidate’ able to provide the condition $p < -\rho/3$. For sure, inflation is a phase of the history of the universe occurring before the era of nucleosynthesis ($t \approx 1$ sec, $T \approx 1$ MeV) during which the light elements abundances were formed. This is because nucleosynthesis is the earliest epoch we have experimental data from and they are in agreement with the predictions of the standard Big-Bang theory. However, the thermal history of the universe before the epoch of nucleosynthesis is unknown.

In order to study the properties of the period of inflation, we assume the extreme condition $p = -\rho$ which considerably simplifies the analysis. A period of the universe during which $p = -\rho$ is called *de Sitter* stage. By inspecting Eqs. (3) and (4), we learn that during the de Sitter phase

$$\begin{aligned}\rho &= \text{constant}, \\ H_I &= \text{constant},\end{aligned}$$

where we have indicated by H_I the value of the Hubble rate during inflation. Correspondingly, solving Eq. (3) gives

$$a = a_i e^{H_I(t-t_i)}, \quad (43)$$

where t_i denotes the time at which inflation starts. Let us now see how such a period of exponential expansion takes care of the shortcomings of the standard Big Bang Theory.⁴

4.1 Inflation and the horizon Problem

During the inflationary (de Sitter) epoch the horizon scale H^{-1} is constant. If inflation lasts long enough, all the physical scales that have left the horizon during the radiation-dominated or matter-dominated phase can re-enter the horizon in the past: this is because such scales are exponentially reduced. As we have seen in the previous section, this explains both the problem of the homogeneity of CMB and the initial condition problem of small cosmological perturbations. Once the physical length is within the horizon, microphysics can act, the universe can be made approximately homogeneous and the *primaeval* inhomogeneities can be created.

Let us see how long inflation must be sustained in order to solve the horizon problem. Let t_i and t_f be, respectively, the time of beginning and end of inflation. We can define the corresponding number of e-foldings N

$$N = \ln [H_I(t_f - t_i)]. \quad (44)$$

A necessary condition to solve the horizon problem is that the largest scale we observe today, the present horizon H_0^{-1} , was reduced during inflation to a value $\lambda_{H_0}(t_i)$ smaller than the value of horizon length H_I^{-1} during inflation. This gives

$$\lambda_{H_0}(t_i) = H_0^{-1} \left(\frac{a_{t_f}}{a_{t_0}} \right) \left(\frac{a_{t_i}}{a_{t_f}} \right) = H_0^{-1} \left(\frac{T_0}{T_f} \right) e^{-N} \lesssim H_I^{-1},$$

⁴ Despite the fact that the growth of the scale factor is exponential and the expansion is *superluminal*, this is not in contradiction with what dictated by relativity. Indeed, it is the spacetime itself which is propagating so fast and not a light signal in it.

where we have neglected for simplicity the short period of matter-domination and we have called T_f the temperature at the end of inflation (to be indentified with the reheating temperature T_{RH} at the beginning of the radiation-dominated phase after inflation, see later). We get

$$N \gtrsim \ln\left(\frac{T_0}{H_0}\right) - \ln\left(\frac{T_f}{H_I}\right) \approx 67 + \ln\left(\frac{T_f}{H_I}\right).$$

Apart from the logarithmic dependence, we obtain $N \gtrsim 70$.

4.2 Inflation and the flatness problem

Inflation solves elegantly the flatness problem. Since during inflation the Hubble rate is constant

$$\Omega - 1 = \frac{k}{a^2 H^2} \propto \frac{1}{a^2}.$$

On the other end the condition (32) tells us that to reproduce a value of $(\Omega_0 - 1)$ of order of unity today the initial value of $(\Omega - 1)$ at the beginning of the radiation-dominated phase must be $|\Omega - 1| \sim 10^{-60}$. Since we identify the beginning of the radiation-dominated phase with the beginning of inflation, we require

$$|\Omega - 1|_{t=t_f} \sim 10^{-60}.$$

During inflation

$$\frac{|\Omega - 1|_{t=t_f}}{|\Omega - 1|_{t=t_i}} = \left(\frac{a_i}{a_f}\right)^2 = e^{-2N}. \quad (45)$$

Taking $|\Omega - 1|_{t=t_i}$ of order unity, it is enough to require that $N \approx 70$ to solve the flatness problem.

1. *Comment:* In the previous section we have written that the flatness problem can be also seen as a fine-tuning problem of one part over 10^{60} . Inflation ameliorates this fine-tuning problem, by explaining a tiny number $\sim 10^{-60}$ with a number N of the order 70.

2. *Comment:* The number $N \simeq 70$ has been obtained requiring that the present-day value of $(\Omega_0 - 1)$ is of order unity. For the expression (45), it is clear that –if the period of inflation lasts longer than 70 e-foldings the present-day value of Ω_0 will be equal to unity with a great precision. One can say that a generic prediction of inflation is that

$$\text{INFLATION} \implies \Omega_0 = 1.$$

This statement, however, must be taken *cum grano salis* and properly specified. Inflation does not change the global geometric properties of the spacetime. If the universe is open or closed, it will always remain flat or closed, independently from inflation. What inflation does is to magnify the radius of curvature R_{curv} defined in Eq. (9) so that locally the universe is flat with a great precision. As we have seen in section 2, the current data on the CMB anisotropies confirm this prediction!

4.3 Inflation and the entropy problem

In the previous section, we have seen that the flatness problem arises because the entropy in a comoving volume is conserved. It is possible, therefore, that the problem could be resolved if the cosmic expansion was non-adiabatic for some finite time interval during the early history of the universe. We need to produce a large amount of entropy $S_U \sim 10^{90}$. Let us postulate that the entropy changed by an amount

$$S_f = Z^3 S_i \quad (46)$$

from the beginning to the end of the inflationary period, where Z is a numerical factor. It is very natural to assume that the total entropy of the universe at the beginning of inflation was of order unity, one particle per horizon. Since, from the end of inflation onwards, the universe expands adiabatically, we have $S_f = S_U$. This gives $Z \sim 10^{30}$. On the other hand, since $S_f \sim (a_f T_f)^3$ and $S_i \sim (a_i T_i)^3$, where T_f and T_i are the temperatures of the universe at the end and at the beginning of inflation, we get

$$\left(\frac{a_f}{a_i}\right) = e^N \approx 10^{30} \left(\frac{T_i}{T_f}\right), \quad (47)$$

which gives again $N \sim 70$ up to the logarithmic factor $\ln\left(\frac{T_i}{T_f}\right)$. We stress again that such a large amount of entropy is not produced during inflation, but during the non-adiabatic phase transition which gives rise to the usual radiation-dominated phase.

4.4 Inflation and the inflaton

In the previous subsections we have described the various advantages of having a period of accelerating phase. The latter required $p < -\rho/3$. Now, we would like to show that this condition can be attained by means of a simple scalar field. We shall call this field the *inflaton* ϕ .

The action of the inflaton field reads

$$S = \int d^4x \sqrt{-g} \mathcal{L} = \int d^4x \sqrt{-g} \left[\frac{1}{2} \partial_\mu \phi \partial^\mu \phi + V(\phi) \right], \quad (48)$$

where $\sqrt{-g} = a^3$ for the FRW metric (1). From the Eulero-Lagrange equations

$$\partial^\mu \frac{\delta(\sqrt{-g} \mathcal{L})}{\delta \partial^\mu \phi} - \frac{\delta(\sqrt{-g} \mathcal{L})}{\delta \phi} = 0, \quad (49)$$

we obtain

$$\ddot{\phi} + 3H\dot{\phi} - \frac{\nabla^2 \phi}{a^2} + V'(\phi) = 0, \quad (50)$$

where $V'(\phi) = (dV(\phi)/d\phi)$. Note, in particular, the appearance of the friction term $3H\dot{\phi}$: a scalar field rolling down its potential suffers a friction due to the expansion of the universe.

We can write the energy-momentum tensor of the scalar field

$$T_{\mu\nu} = \partial_\mu \phi \partial_\nu \phi - g_{\mu\nu} \mathcal{L}.$$

The corresponding energy density ρ_ϕ and pressure density p_ϕ are

$$T_{00} = \rho_\phi = \frac{\dot{\phi}^2}{2} + V(\phi) + \frac{(\nabla\phi)^2}{2a^2}, \quad (51)$$

$$T_{ii} = p_\phi = \frac{\dot{\phi}^2}{2} - V(\phi) - \frac{(\nabla\phi)^2}{6a^2}. \quad (52)$$

Notice that, if the gradient term were dominant, we would obtain $p_\phi = -\frac{\rho_\phi}{3}$, not enough to drive inflation. We can now split the inflaton field in

$$\phi(t) = \phi_0(t) + \delta\phi(\mathbf{x}, t),$$

where ϕ_0 is the ‘classical’ (infinite wavelength) field, that is the expectation value of the inflaton field on the initial isotropic and homogeneous state, while $\delta\phi(\mathbf{x}, t)$ represents the quantum fluctuations around ϕ_0 . In this section, we will be only concerned with the evolution of the classical field ϕ_0 . The next section will be devoted to the crucial issue of the evolution of quantum perturbations during inflation. This separation is justified by the fact that quantum fluctuations are much smaller than the classical value and therefore negligible when looking at the classical evolution. To not be overwhelmed by the notation, we will keep indicating from now on the classical value of the inflaton field by ϕ . The energy-momentum tensor becomes

$$T_{00} = \rho_\phi = \frac{\dot{\phi}^2}{2} + V(\phi) \quad (53)$$

$$T_{ii} = p_\phi = \frac{\dot{\phi}^2}{2} - V(\phi). \quad (54)$$

If

$$V(\phi) \gg \dot{\phi}^2$$

we obtain the following condition

$$p_\phi \simeq -\rho_\phi$$

From this simple calculation, we realize that a scalar field whose energy is dominant in the universe and whose potential energy dominates over the kinetic term gives inflation! Inflation is driven by the vacuum energy of the inflaton field.

4.5 Slow-roll conditions

Let us now quantify better under which circumstances a scalar field may give rise to a period of inflation. The equation of motion of the field is

$$\ddot{\phi} + 3H\dot{\phi} + V'(\phi) = 0 \quad (55)$$

If we require that $\dot{\phi}^2 \ll V(\phi)$, the scalar field is slowly rolling down its potential. This is the reason why such a period is called *slow-roll*. We may also expect that – being the potential

flat $-\ddot{\phi}$ is negligible as well. We will assume that this is true and we will quantify this condition soon. The FRW equation (3) becomes

$$H^2 \simeq \frac{8\pi G}{3} V(\phi), \quad (56)$$

where we have assumed that the inflaton field dominates the energy density of the universe. The new equation of motion becomes

$$3H\dot{\phi} = -V'(\phi) \quad (57)$$

which gives $\dot{\phi}$ as a function of $V'(\phi)$. Using Eq. (57) slow-roll conditions then require

$$\dot{\phi}^2 \ll V(\phi) \implies \frac{(V')^2}{V} \ll H^2$$

and

$$\ddot{\phi} \ll 3H\dot{\phi} \implies V'' \ll H^2.$$

It is now useful to define the slow-roll parameters, ϵ and η in the following way

$$\begin{aligned} \epsilon &= -\frac{\dot{H}}{H^2} = 4\pi G \frac{\dot{\phi}^2}{H^2} = \frac{1}{16\pi G} \left(\frac{V'}{V} \right)^2, \\ \eta &= \frac{1}{8\pi G} \left(\frac{V''}{V} \right) = \frac{1}{3} \frac{V''}{H^2}, \\ \delta &= \eta - \epsilon = -\frac{\ddot{\phi}}{H\dot{\phi}}. \end{aligned}$$

It might be useful to have the same parameters expressed in terms of conformal time

$$\begin{aligned} \epsilon &= 1 - \frac{\mathcal{H}'}{\mathcal{H}^2} = 4\pi G \frac{\phi'^2}{\mathcal{H}^2} \\ \delta &= \eta - \epsilon = 1 - \frac{\phi''}{\mathcal{H}\phi'}. \end{aligned}$$

The parameter ϵ quantifies how much the Hubble rate H changes with time during inflation. Notice that, since

$$\frac{\ddot{a}}{a} = \dot{H} + H^2 = (1 - \epsilon) H^2,$$

inflation can be attained only if $\epsilon < 1$:

$$\text{INFLATION} \iff \epsilon < 1.$$

As soon as this condition fails, inflation ends. In general, slow-roll inflation is attained if $\epsilon \ll 1$ and $|\eta| \ll 1$. During inflation the slow-roll parameters ϵ and η can be considered to be approximately constant since the potential $V(\phi)$ is very flat.

Comment: In the following, we will work at *first-order* perturbation in the slow-roll parameters, that is we will take only the first power of them. Since, using their definition, it is easy to see that $\dot{\epsilon}, \dot{\eta} = \mathcal{O}(\epsilon^2, \eta^2)$, this amounts to saying that we will treat the slow-roll parameters as constant in time.

Within these approximations, it is easy to compute the number of e-foldings between the beginning and the end of inflation. If we indicate by ϕ_i and ϕ_f the values of the inflaton field at the beginning and at the end of inflation, respectively, we have that the *total* number of e-foldings is

$$\begin{aligned}
N &\equiv \int_{t_i}^{t_f} H dt \\
&\simeq H \int_{\phi_i}^{\phi_f} \frac{d\phi}{\dot{\phi}} \\
&\simeq -3H^2 \int_{\phi_i}^{\phi_f} \frac{d\phi}{V'} \\
&\simeq -8\pi G \int_{\phi_i}^{\phi_f} \frac{V}{V'} d\phi.
\end{aligned} \tag{58}$$

We may also compute the number of e-foldings ΔN which are left to go to the end of inflation

$$\Delta N \simeq 8\pi G \int_{\phi_f}^{\phi_{\Delta N}} \frac{V}{V'} d\phi, \tag{59}$$

where $\phi_{\Delta N}$ is the value of the inflaton field when there are ΔN e-foldings to the end of inflation.

1. Comment: According to the criterion given in subsection 2.4, a given scale length $\lambda = a/k$ leaves the horizon when $k = aH_k$ where H_k is the value of the Hubble rate at that time. One can compute easily the rate of change of H_k^2 as a function of k

$$\frac{d \ln H_k^2}{d \ln k} = \left(\frac{d \ln H_k^2}{dt} \right) \left(\frac{dt}{d \ln a} \right) \left(\frac{d \ln a}{d \ln k} \right) = 2 \frac{\dot{H}}{H} \times \frac{1}{H} \times 1 = 2 \frac{\dot{H}}{H^2} = -2\epsilon. \tag{60}$$

2. Comment: Take a given physical scale λ today which crossed the horizon scale during inflation. This happened when

$$\lambda \left(\frac{a_f}{a_0} \right) e^{-\Delta N_\lambda} = \lambda \left(\frac{T_0}{T_f} \right) e^{-\Delta N_\lambda} = H_I^{-1}$$

where ΔN_λ indicates the number of e-foldings from the time the scale crossed the horizon during inflation and the end of inflation. This relation gives a way to determine the number of e-foldings to the end of inflation corresponding to a given scale

$$\Delta N_\lambda \simeq 65 + \ln \left(\frac{\lambda}{3000 \text{ Mpc}} \right) + 2 \ln \left(\frac{V^{1/4}}{10^{14} \text{ GeV}} \right) - \ln \left(\frac{T_f}{10^{10} \text{ GeV}} \right).$$

Scales relevant for the CMB anisotropies correspond to $\Delta N \sim 60$.

4.6 The last stage of inflation and reheating

Inflation ended when the potential energy associated with the inflaton field became smaller than the kinetic energy of the field. By that time, any pre-inflation entropy in the universe had been inflated away, and the energy of the universe was entirely in the form of coherent oscillations of the inflaton condensate around the minimum of its potential. The universe may be said to be frozen after the end of inflation. We know that somehow the low-entropy cold universe dominated by the energy of coherent motion of the ϕ field must be transformed into a high-entropy hot universe dominated by radiation. The process by which the energy of the inflaton field is transferred from the inflaton field to radiation has been dubbed *reheating*. In the old theory of reheating [21, 1], the simplest way to envision this process is if the comoving energy density in the zero mode of the inflaton decays into normal particles, which then scatter and thermalize to form a thermal background. It is usually assumed that the decay width of this process is the same as the decay width of a free inflaton field.

Of particular interest is a quantity known usually as the reheat temperature, denoted as T_{RH} ⁵. The reheat temperature is calculated by assuming an instantaneous conversion of the energy density in the inflaton field into radiation when the decay width of the inflaton energy, Γ_ϕ , is equal to H , the expansion rate of the universe.

The reheat temperature is calculated quite easily. After inflation the inflaton field executes coherent oscillations about the minimum of the potential. Averaged over several oscillations, the coherent oscillation energy density redshifts as matter: $\rho_\phi \propto a^{-3}$, where a is the Robertson–Walker scale factor. If we denote as ρ_I and a_I the total inflaton energy density and the scale factor at the initiation of coherent oscillations, then the Hubble expansion rate as a function of a is

$$H^2(a) = \frac{8\pi}{3} \frac{\rho_I}{m_{\text{Pl}}^2} \left(\frac{a_I}{a} \right)^3. \quad (61)$$

Equating $H(a)$ and Γ_ϕ leads to an expression for a_I/a . Now if we assume that all available coherent energy density is instantaneously converted into radiation at this value of a_I/a , we can find the reheat temperature by setting the coherent energy density, $\rho_\phi = \rho_I(a_I/a)^3$, equal to the radiation energy density, $\rho_R = (\pi^2/30)g_*T_{RH}^4$, where g_* is the effective number of relativistic degrees of freedom at temperature T_{RH} . The result is

$$T_{RH} = \left(\frac{90}{8\pi^3 g_*} \right)^{1/4} \sqrt{\Gamma_\phi m_{\text{Pl}}} = 0.2 \left(\frac{200}{g_*} \right)^{1/4} \sqrt{\Gamma_\phi m_{\text{Pl}}}. \quad (62)$$

⁵So far, we have indicated it with T_f .

In some models of inflation reheating can be anticipated by a period of preheating [35] when the the classical inflaton field very rapidly (explosively) decays into ϕ -particles or into other bosons due to broad parametric resonance. This stage cannot be described by the standard elementary approach to reheating based on perturbation theory. The bosons produced at this stage further decay into other particles, which eventually become thermalized.

4.7 A brief survey of inflationary models

Even restricting ourselves to a simple single-field inflation scenario, the number of models available to choose from is large [48]. It is convenient to define a general classification scheme, or “zoology” for models of inflation. We divide models into three general types [20]: *large-field*, *small-field*, and *hybrid*, with a fourth classification. A generic single-field potential can be characterized by two independent mass scales: a “height” Λ^4 , corresponding to the vacuum energy density during inflation, and a “width” μ , corresponding to the change in the field value $\Delta\phi$ during inflation:

$$V(\phi) = \Lambda^4 f\left(\frac{\phi}{\mu}\right). \quad (63)$$

Different models have different forms for the function f . Let us now briefly describe the different class of models.

4.7.1 Large-field models

Large-field models are potentials typical of the “chaotic” inflation scenario[43], in which the scalar field is displaced from the minimum of the potential by an amount usually of order the Planck mass. Such models are characterized by $V''(\phi) > 0$, and $-\epsilon < \delta \leq \epsilon$. The generic large-field potentials we consider are polynomial potentials $V(\phi) = \Lambda^4 (\phi/\mu)^p$, and exponential potentials, $V(\phi) = \Lambda^4 \exp(\phi/\mu)$. In the chaotic inflation scenario, it is assumed that the universe emerged from a quantum gravitational state with an energy density comparable to that of the Planck density. This implies that $V(\phi) \approx m_{\text{Pl}}^4$ and results in a large friction term in the Friedmann equation (180). Consequently, the inflaton will slowly roll down its potential. The condition for inflation is therefore satisfied and the scale factor grows as

$$a(t) = a_i e^{\left(\int_{t_i}^t dt' H(t')\right)}. \quad (64)$$

The simplest chaotic inflation model is that of a free field with a quadratic potential, $V(\phi) = m^2 \phi^2/2$, where m represents the mass of the inflaton. During inflation the scale factor grows as

$$a(t) = a_i e^{2\pi(\phi_i^2 - \phi^2(t))} \quad (65)$$

and inflation ends when $\phi = \mathcal{O}(1) m_{\text{Pl}}$. If inflation begins when $V(\phi_i) \approx m_{\text{Pl}}^4$, the scale factor grows by a factor $\exp(4\pi m_{\text{Pl}}^2/m^2)$ before the inflaton reaches the minimum of its potential. We will later show that the mass of the field should be $m \approx 10^{-6} m_{\text{Pl}}$ if the microwave background constraints are to be satisfied. This implies that the volume of the

universe will increase by a factor of $Z^3 \approx 10^{3 \times 10^{12}}$ and this is more than enough inflation to solve the problems of the hot big bang model.

In the chaotic inflationary scenarios, the present-day universe is only a small portion of the universe which suffered inflation! Notice also that the typical values of the inflaton field during inflation are of the order of m_{Pl} , giving rise to the possibility of testing planckian physics [17].

4.7.2 Small-field models

Small-field models are the type of potentials that arise naturally from spontaneous symmetry breaking (such as the original models of “new” inflation [42, 3]) and from pseudo Nambu-Goldstone modes (natural inflation[24]). The field starts from near an unstable equilibrium (taken to be at the origin) and rolls down the potential to a stable minimum. Small-field models are characterized by $V''(\phi) < 0$ and $\eta < -\epsilon$. Typically ϵ is close to zero. The generic small-field potentials we consider are of the form $V(\phi) = \Lambda^4 [1 - (\phi/\mu)^p]$, which can be viewed as a lowest-order Taylor expansion of an arbitrary potential about the origin. See, for instance, Ref. [19].

4.7.3 Hybrid models

The hybrid scenario[44, 45, 16] frequently appears in models which incorporate inflation into supersymmetry [57] and supergravity [46]. In a typical hybrid inflation model, the scalar field responsible for inflation evolves toward a minimum with nonzero vacuum energy. The end of inflation arises as a result of instability in a second field. Such models are characterized by $V''(\phi) > 0$ and $0 < \epsilon < \delta$. We consider generic potentials for hybrid inflation of the form $V(\phi) = \Lambda^4 [1 + (\phi/\mu)^p]$. The field value at the end of inflation is determined by some other physics, so there is a second free parameter characterizing the models. This enumeration of models is certainly not exhaustive. There are a number of single-field models that do not fit well into this scheme, for example logarithmic potentials $V(\phi) \propto \ln(\phi)$ typical of supersymmetry [48, 28, 13, 22, 47, 58, 23, 34]. Another example is potentials with negative powers of the scalar field $V(\phi) \propto \phi^{-p}$ used in intermediate inflation [8] and dynamical supersymmetric inflation [31, 32]. Both of these cases require an auxiliary field to end inflation and are more properly categorized as hybrid models, but fall into the small-field class. However, the three classes categorized by the relationship between the slow-roll parameters as $-\epsilon < \delta \leq \epsilon$ (large-field), $\delta \leq -\epsilon$ (small-field) and $0 < \epsilon < \delta$ (hybrid) seems to be good enough for comparing theoretical expectations with experimental data.

5 Inflation and the cosmological perturbations

As we have seen in the previous section, the early universe was made very nearly uniform by a primordial inflationary stage. However, the important caveat in that statement is the word ‘nearly’. Our current understanding of the origin of structure in the universe is that it originated from small ‘seed’ perturbations, which over time grew to become all of the structure we observe. Once the universe becomes matter dominated (around 1000 yrs after

the bang) primeval density inhomogeneities ($\delta\rho/\rho \sim 10^{-5}$) are amplified by gravity and grow into the structure we see today [54]. The fact that a fluid of self-gravitating particles is unstable to the growth of small inhomogeneities was first pointed out by Jeans and is known as the Jeans instability. Furthermore, the existence of these inhomogeneities was confirmed by the COBE discovery of CMB anisotropies; the temperature anisotropies detected almost certainly owe their existence to primeval density inhomogeneities, since, as we have seen, causality precludes microphysical processes from producing anisotropies on angular scales larger than about 1° , the angular size of the horizon at last-scattering.

The growth of small matter inhomogeneities of wavelength smaller than the Hubble scale ($\lambda \lesssim H^{-1}$) is governed by a Newtonian equation:

$$\ddot{\delta}_{\mathbf{k}} + 2H\dot{\delta}_{\mathbf{k}} + v_s^2 \frac{k^2}{a^2} \delta_{\mathbf{k}} = 4\pi G \rho_M \delta_{\mathbf{k}}, \quad (66)$$

where $v_s^2 = \partial p / \partial \rho_M$ is the square of the sound speed and we have expanded the perturbation to the matter density in plane waves

$$\frac{\delta\rho_M(\mathbf{x}, t)}{\rho_M} = \frac{1}{(2\pi)^3} \int d^3k \delta_{\mathbf{k}}(t) e^{-i\mathbf{k}\cdot\mathbf{x}}. \quad (67)$$

Competition between the pressure term and the gravity term on the rhs of Eq. (66) determines whether or not pressure can counteract gravity: perturbations with wavenumber larger than the Jeans wavenumber, $k_J^2 = 4\pi G a^2 \rho_M / v_s^2$, are Jeans stable and just oscillate; perturbations with smaller wavenumber are Jeans unstable and can grow.

Let us discuss solutions to this equation under different circumstances. First, consider the Jeans problem, evolution of perturbations in a static fluid, *i.e.*, $H = 0$. In this case Jeans unstable perturbations grow exponentially, $\delta_{\mathbf{k}} \propto \exp(t/\tau)$ where $\tau = 1/\sqrt{4\pi G \rho_M}$. Next, consider the growth of Jeans unstable perturbations in a matter-dominated universe, *i.e.*, $H^2 = 8\pi G \rho_M / 3$ and $a \propto t^{2/3}$. Because the expansion tends to “pull particles away from one another,” the growth is only power law, $\delta_{\mathbf{k}} \propto t^{2/3}$; *i.e.*, at the same rate as the scale factor. Finally, consider a radiation-dominated universe. In this case, the expansion is so rapid that matter perturbations grow very slowly, as $\ln a$ in radiation-dominated epoch. Therefore, perturbations may grow only in a matter-dominated period. Once a perturbation reaches an overdensity of order unity or larger it “separates” from the expansion *—i.e.*, becomes its own self-gravitating system and ceases to expand any further. In the process of virial relaxation, its size decreases by a factor of two—density increases by a factor of 8; thereafter, its density contrast grows as a^3 since the average matter density is decreasing as a^{-3} , though smaller scales could become Jeans unstable and collapse further to form smaller objects of higher density.

In order for structure formation to occur via gravitational instability, there must have been small preexisting fluctuations on physical length scales when they crossed the Hubble radius in the radiation-dominated and matter-dominated eras. In the standard Big-Bang model these small perturbations have to be put in by hand, because it is impossible to produce fluctuations on any length scale while it is larger than the horizon. Since the goal of cosmology is to understand the universe on the basis of physical laws, this appeal to initial conditions is unsatisfactory. The challenge is therefore to give an explanation to the small seed perturbations which allow the gravitational growth of the matter perturbations.

Our best guess for the origin of these perturbations is quantum fluctuations during an inflationary era in the early universe. Although originally introduced as a possible solution to the cosmological conundrums such as the horizon, flatness and entropy problems, by far the most useful property of inflation is that it generates spectra of both density perturbations and gravitational waves. These perturbations extend from extremely short scales to scales considerably in excess of the size of the observable universe.

During inflation the scale factor grows quasi-exponentially, while the Hubble radius remains almost constant. Consequently the wavelength of a quantum fluctuation – either in the scalar field whose potential energy drives inflation or in the graviton field – soon exceeds the Hubble radius. The amplitude of the fluctuation therefore becomes ‘frozen in’. This is quantum mechanics in action at macroscopic scales!

According to quantum field theory, empty space is not entirely empty. It is filled with quantum fluctuations of all types of physical fields. The fluctuations can be regarded as waves of physical fields with all possible wavelengths, moving in all possible directions. If the values of these fields, averaged over some macroscopically large time, vanish then the space filled with these fields seems to us empty and can be called the vacuum.

In the exponentially expanding universe the vacuum structure is much more complicated. The wavelengths of all vacuum fluctuations of the inflaton field ϕ grow exponentially in the expanding universe. When the wavelength of any particular fluctuation becomes greater than H^{-1} , this fluctuation stops propagating, and its amplitude freezes at some nonzero value $\delta\phi$ because of the large friction term $3H\dot{\phi}$ in the equation of motion of the field ϕ . The amplitude of this fluctuation then remains almost unchanged for a very long time, whereas its wavelength grows exponentially. Therefore, the appearance of such frozen fluctuation is equivalent to the appearance of a classical field $\delta\phi$ that does not vanish after having averaged over some macroscopic interval of time. Because the vacuum contains fluctuations of all possible wavelength, inflation leads to the creation of more and more new perturbations of the classical field with wavelength larger than the horizon scale.

Once inflation has ended, however, the Hubble radius increases faster than the scale factor, so the fluctuations eventually reenter the Hubble radius during the radiation- or matter-dominated eras. The fluctuations that exit around 60 e -foldings or so before reheating reenter with physical wavelengths in the range accessible to cosmological observations. These spectra provide a distinctive signature of inflation. They can be measured in a variety of different ways including the analysis of microwave background anisotropies.

The physical processes which give rise to the structures we observe today are well-explained in Fig. 8. Quantum fluctuations of the inflaton field are generated during inflation. Since gravity talks to any component of the universe, small fluctuations of the inflaton field are intimately related to fluctuations of the spacetime metric, giving rise to perturbations of the curvature $\mathcal{R}$ (which will be defined in the following; the reader may loosely think of it as a gravitational potential). The wavelengths λ of these perturbations grow exponentially and leave soon the horizon when $\lambda > R_H$. On superhorizon scales, curvature fluctuations are frozen in and may be considered as classical. Finally, when the wavelength of these fluctuations reenters the horizon, at some radiation- or matter-dominated epoch, the curvature (gravitational potential) perturbations of the spacetime give rise to matter (and temperature) perturbations $\delta\rho$ via the Poisson equation. These fluctuations will then start growing giving rise to the structures we observe today.

In summary, two are the key ingredients for understanding the observed structures in the

universe within the inflationary scenario:

- Quantum fluctuations of the inflaton field are excited during inflation and stretched to cosmological scales. At the same time, being the inflaton fluctuations connected to the metric perturbations through Einstein's equations, ripples on the metric are also excited and stretched to cosmological scales.
- Gravity acts a messenger since it communicates to baryons and photons the small seed perturbations once a given wavelength becomes smaller than the horizon scale after inflation.

Let us now see how quantum fluctuations are generated during inflation. We will proceed by steps. First, we will consider the simplest problem of studying the quantum fluctuations of a generic scalar field during inflation: we will learn how perturbations evolve as a function of time and compute their spectrum. Then – since a satisfactory description of the generation of quantum fluctuations have to take both the inflaton and the metric perturbations into account – we will study the system composed by quantum fluctuations of the inflaton field and quantum fluctuations of the metric.

6 Quantum fluctuations of a generic massless scalar field during inflation

Let us first see how the fluctuations of a generic scalar field χ , which is *not* the inflaton field, behave during inflation. To warm up we first consider a de Sitter epoch during which the Hubble rate is constant.

6.1 Quantum fluctuations of a generic massless scalar field during a de Sitter stage

We assume this field to be massless. The massive case will be analyzed in the next subsection. Expanding the scalar field χ in Fourier modes

$$\delta\chi(\mathbf{x}, t) = \int \frac{d^3\mathbf{k}}{(2\pi)^{3/2}} e^{i\mathbf{k}\cdot\mathbf{x}} \delta\chi_{\mathbf{k}}(t),$$

we can write the equation for the fluctuations as

$$\delta\ddot{\chi}_{\mathbf{k}} + 3H \delta\dot{\chi}_{\mathbf{k}} + \frac{k^2}{a^2} \delta\chi_{\mathbf{k}} = 0. \quad (68)$$

Let us study the qualitative behaviour of the solution to Eq. (68).

- For wavelengths within the horizon, $\lambda \ll H^{-1}$, the corresponding wavenumber satisfies the relation $k \gg aH$. In this regime, we can neglect the friction term $3H \delta\dot{\chi}_{\mathbf{k}}$ and Eq. (68) reduces to

$$\delta\ddot{\chi}_{\mathbf{k}} + \frac{k^2}{a^2} \delta\chi_{\mathbf{k}} = 0, \quad (69)$$

which is – basically – the equation of motion of an harmonic oscillator. Of course, the frequency term k^2/a^2 depends upon time because the scale factor a grows exponentially. On the qualitative level, however, one expects that when the wavelength of the fluctuation is within the horizon, the fluctuation oscillates.

- For wavelengths above the horizon, $\lambda \gg H^{-1}$, the corresponding wavenumber satisfies the relation $k \ll aH$ and the term k^2/a^2 can be safely neglected. Eq. (68) reduces to

$$\delta\ddot{\chi}_{\mathbf{k}} + 3H \delta\dot{\chi}_{\mathbf{k}} = 0, \quad (70)$$

which tells us that on superhorizon scales $\delta\chi_{\mathbf{k}}$ remains constant.

We have therefore the following picture: take a given fluctuation whose initial wavelength $\lambda \sim a/k$ is within the horizon. The fluctuations oscillates till the wavelength becomes of the order of the horizon scale. When the wavelength crosses the horizon, the fluctuation ceases to oscillate and gets frozen in.

Let us now study the evolution of the fluctuation in a more quantitative way. To do so, we perform the following redefinition

$$\delta\chi_{\mathbf{k}} = \frac{\delta\sigma_{\mathbf{k}}}{a}$$

and we work in conformal time $d\tau = dt/a$. For the time being, we solve the problem for a pure de Sitter expansion and we take the scale factor exponentially growing as $a \sim e^{Ht}$; the corresponding conformal factor reads (after choosing properly the integration constants)

$$a(\tau) = -\frac{1}{H\tau} \quad (\tau < 0).$$

In the following we will also solve the problem in the case of quasi de Sitter expansion. The beginning of inflation coincides with some initial time $\tau_i \ll 0$. Using the set of rules (15), we find that Eq. (68) becomes

$$\delta\sigma_{\mathbf{k}}'' + \left(k^2 - \frac{a''}{a}\right) \delta\sigma_{\mathbf{k}} = 0. \quad (71)$$

We obtain an equation which is very ‘close’ to the equation for a Klein-Gordon scalar field in flat spacetime, the only difference being a negative time-dependent mass term $-a''/a = -2/\tau^2$. Eq. (71) can be obtained from an action of the type

$$\delta S_{\mathbf{k}} = \int d\tau \left[\frac{1}{2} \delta\sigma_{\mathbf{k}}'^2 - \frac{1}{2} \left(k^2 - \frac{a''}{a}\right) \delta\sigma_{\mathbf{k}}^2 \right], \quad (72)$$

which is the canonical action for a simple harmonic oscillator with canonical commutation relations $\delta\sigma_{\mathbf{k}}^* \delta\sigma_{\mathbf{k}}' - \delta\sigma_{\mathbf{k}} \delta\sigma_{\mathbf{k}}^{*'} = -i$.

Let us study the behaviour of this equation on subhorizon and superhorizon scales. Since

$$\frac{k}{aH} = -k\tau,$$

on subhorizon scales $k^2 \gg a''/a$ Eq. (71) reduces to

$$\delta\sigma_{\mathbf{k}}'' + k^2 \delta\sigma_{\mathbf{k}} = 0,$$

whose solution is a plane wave

$$\delta\sigma_{\mathbf{k}} = \frac{e^{-ik\tau}}{\sqrt{2k}} \quad (k \gg aH). \quad (73)$$

We find again that fluctuations with wavelength within the horizon oscillate exactly like in flat spacetime. This does not come as a surprise. In the ultraviolet regime, that is for wavelengths much smaller than the horizon scale, one expects that approximating the spacetime as flat is a good approximation.

On superhorizon scales, $k^2 \ll a''/a$ Eq. (71) reduces to

$$\delta\sigma_{\mathbf{k}}'' - \frac{a''}{a} \delta\sigma_{\mathbf{k}} = 0,$$

which is satisfied by

$$\delta\sigma_{\mathbf{k}} = B(k) a \quad (k \ll aH). \quad (74)$$

where $B(k)$ is a constant of integration. Roughly matching the (absolute values of the) solutions (73) and (74) at $k = aH$ ($-k\tau = 1$), we can determine the (absolute value of the) constant $B(k)$

$$|B(k)| a = \frac{1}{\sqrt{2k}} \implies |B(k)| = \frac{1}{a\sqrt{2k}} = \frac{H}{\sqrt{2k^3}}.$$

Going back to the original variable $\delta\chi_{\mathbf{k}}$, we obtain that the quantum fluctuation of the χ field on superhorizon scales is constant and approximately equal to

$$|\delta\chi_{\mathbf{k}}| \simeq \frac{H}{\sqrt{2k^3}} \quad (\text{ON SUPERHORIZON SCALES})$$

In fact we can do much better, since Eq. (71) has an *exact* solution:

$$\delta\sigma_{\mathbf{k}} = \frac{e^{-ik\tau}}{\sqrt{2k}} \left(1 + \frac{i}{k\tau} \right). \quad (75)$$

This solution reproduces all what we have found by qualitative arguments in the two extreme regimes $k \ll aH$ and $k \gg aH$. The reason why we have performed the matching procedure is to show that the latter can be very useful to determine the behaviour of the solution on superhorizon scales when the exact solution is not known.

6.2 Quantum fluctuations of a generic massive scalar field during a de Sitter stage

So far, we have solved the equation for the quantum perturbations of a generic massless field, that is neglecting the mass squared term m_χ^2 . Let us now discuss the solution when such a mass term is present. Eq. (71) becomes

$$\delta\sigma_{\mathbf{k}}'' + [k^2 + M^2(\tau)] \delta\sigma_{\mathbf{k}} = 0, \quad (76)$$

where

$$M^2(\tau) = (m_\chi^2 - 2H^2) a^2(\tau) = \frac{1}{\tau^2} \left(\frac{m_\chi^2}{H^2} - 2 \right).$$

Eq. (76) can be recast in the form

$$\delta\sigma_{\mathbf{k}}'' + \left[k^2 - \frac{1}{\tau^2} \left(\nu_\chi^2 - \frac{1}{4} \right) \right] \delta\sigma_{\mathbf{k}} = 0, \quad (77)$$

where

$$\nu_\chi^2 = \left(\frac{9}{4} - \frac{m_\chi^2}{H^2} \right). \quad (78)$$

The generic solution to Eq. (76) for ν_χ *real* is

$$\delta\sigma_{\mathbf{k}} = \sqrt{-\tau} \left[c_1(k) H_{\nu_\chi}^{(1)}(-k\tau) + c_2(k) H_{\nu_\chi}^{(2)}(-k\tau) \right],$$

where $H_{\nu_\chi}^{(1)}$ and $H_{\nu_\chi}^{(2)}$ are the Hankel's functions of the first and second kind, respectively. If we impose that in the ultraviolet regime $k \gg aH$ ($-k\tau \gg 1$) the solution matches the plane-wave solution $e^{-ik\tau}/\sqrt{2k}$ that we expect in flat spacetime and knowing that

$$H_{\nu_\chi}^{(1)}(x \gg 1) \sim \sqrt{\frac{2}{\pi x}} e^{i(x - \frac{\pi}{2}\nu_\chi - \frac{\pi}{4})}, \quad H_{\nu_\chi}^{(2)}(x \gg 1) \sim \sqrt{\frac{2}{\pi x}} e^{-i(x - \frac{\pi}{2}\nu_\chi - \frac{\pi}{4})},$$

we set $c_2(k) = 0$ and $c_1(k) = \frac{\sqrt{\pi}}{2} e^{i(\nu_\chi + \frac{1}{2})\frac{\pi}{2}}$. The exact solution becomes

$$\delta\sigma_{\mathbf{k}} = \frac{\sqrt{\pi}}{2} e^{i(\nu_\chi + \frac{1}{2})\frac{\pi}{2}} \sqrt{-\tau} H_{\nu_\chi}^{(1)}(-k\tau). \quad (79)$$

On superhorizon scales, since $H_{\nu_\chi}^{(1)}(x \ll 1) \sim \sqrt{2/\pi} e^{-i\frac{\pi}{2}} 2^{\nu_\chi - \frac{3}{2}} (\Gamma(\nu_\chi)/\Gamma(3/2)) x^{-\nu_\chi}$, the fluctuation (79) becomes

$$\delta\sigma_{\mathbf{k}} = e^{i(\nu_\chi - \frac{1}{2})\frac{\pi}{2}} 2^{(\nu_\chi - \frac{3}{2})} \frac{\Gamma(\nu_\chi)}{\Gamma(3/2)} \frac{1}{\sqrt{2k}} (-k\tau)^{\frac{1}{2} - \nu_\chi}.$$

Going back to the old variable $\delta\chi_{\mathbf{k}}$, we find that on superhorizon scales, the fluctuation with nonvanishing mass is not exactly constant, but it acquires a tiny dependence upon the time

$$|\delta\chi_{\mathbf{k}}| \simeq \frac{H}{\sqrt{2k^3}} \left(\frac{k}{aH} \right)^{\frac{3}{2}-\nu_\chi} \quad (\text{ON SUPERHORIZON SCALES})$$

If we now define, in analogy with the definition of the slow roll parameters η and ϵ for the inflaton field, the parameter $\eta_\chi = (m_\chi^2/3H^2) \ll 1$, one finds

$$\frac{3}{2} - \nu_\chi \simeq \eta_\chi. \quad (80)$$

It is instructive to analyze the case in which ν_χ is *imaginary*, that is $m_\chi/H > 3/2$. In such a case, we define $\tilde{\nu} = i\nu$. In superhorizon scales, performing the same steps we have done for the case of ν_χ real, we find

$$|\delta\chi_{\mathbf{k}}|^2 = \frac{\pi}{4} \frac{e^{-\pi\tilde{\nu}}}{a^2} \frac{1}{aH} \left[\frac{(1+\cot(\pi\tilde{\nu}))\sinh(\pi\tilde{\nu})}{\pi\tilde{\nu}} + \frac{\tilde{\nu}}{\pi\sinh(\pi\tilde{\nu})} + 2 \operatorname{Re} \left(i \left(\cos \left(2\tilde{\nu} \ln \left(\frac{k\eta}{2} \right) \right) + i \sin \left(2\tilde{\nu} \ln \left(\frac{k\tau}{2} \right) \right) \right) \frac{1-\cot(\pi\tilde{\nu})}{\Gamma(1+i\tilde{\nu})} \frac{\Gamma(-i\tilde{\nu})}{\pi} \right) \right] \left(\frac{k}{aH} \right)^{n-1}. \quad (81)$$

In the limit of long wavelengths, the highly oscillating term which appears in the real part can be neglected because its average on k is 0. The resulting power spectrum is the following

$$\begin{aligned} \mathcal{P}_{\delta\chi}(k) &= \frac{\pi}{4} e^{-\pi\tilde{\nu}} \left(\frac{H}{2\pi} \right)^2 \left(\frac{(1+\cot(\pi\tilde{\nu}))\sinh(\pi\tilde{\nu})}{\pi\tilde{\nu}} + \frac{\tilde{\nu}}{\pi\sinh(\pi\tilde{\nu})} \right) \\ &\times \left(\frac{k}{aH} \right)^{n-1} \simeq \left(\frac{H}{2\pi} \right)^2 \left(\frac{H}{m_\chi} \right) \left(\frac{k}{aH} \right)^3. \end{aligned} \quad (82)$$

Therefore, for very massive scalar fields, $m_\chi > 3H/2$, the power spectrum has an amplitude which is suppressed by the ratio (H/m_χ) and the spectrum falls down rapidly at large wavelengths k^{-1} as k^3 .

6.3 Quantum to classical transition

We have previously said that the quantum fluctuations can be regarded as classical when their corresponding wavelengths cross the horizon. To better motivate this statement, we should compute the number of particles $n_{\mathbf{k}}$ per wavenumber $\mathbf{k}$ on superhorizon scales and check that it is indeed much larger than unity, $n_{\mathbf{k}} \gg 1$ (in this limit one can neglect the “quantum” factor $1/2$ in the Hamiltonian $H_{\mathbf{k}} = \omega_{\mathbf{k}} \left(n_{\mathbf{k}} + \frac{1}{2} \right)$ where $\omega_{\mathbf{k}}$ is the energy eigenvalue). If so, the fluctuation can be regarded as classical. The number of particles $n_{\mathbf{k}}$ can be estimated to be of the order of $H_{\mathbf{k}}/\omega_{\mathbf{k}}$, where $H_{\mathbf{k}}$ is the Hamiltonian corresponding to the action

$$\delta S_{\mathbf{k}} = \int d\tau \left[\frac{1}{2} \delta\sigma_{\mathbf{k}}'^2 + \frac{1}{2} \left(k^2 - M^2(\tau) \right) \delta\sigma_{\mathbf{k}}^2 \right], \quad (83)$$

One obtains on superhorizon scales

$$n_{\mathbf{k}} \simeq \frac{M^2(\tau) |\delta\chi_{\mathbf{k}}|^2}{\omega_{\mathbf{k}}} \sim \left(\frac{k}{aH} \right)^{-3} \gg 1,$$

which confirms that fluctuations on superhorizon scales may be indeed considered as classical.

6.4 The power spectrum

Let us define now the power spectrum, a useful quantity to characterize the properties of the perturbations. For a generic quantity $g(\mathbf{x}, t)$, which can be expanded in Fourier space as

$$g(\mathbf{x}, t) = \int \frac{d^3\mathbf{k}}{(2\pi)^{3/2}} e^{i\mathbf{k}\cdot\mathbf{x}} g_{\mathbf{k}}(t),$$

the power spectrum can be defined as

$$\langle 0 | g_{\mathbf{k}_1}^* g_{\mathbf{k}_2} | 0 \rangle \equiv \delta^{(3)}(\mathbf{k}_1 - \mathbf{k}_2) \frac{2\pi^2}{k^3} \mathcal{P}_g(k), \quad (84)$$

where $|0\rangle$ is the vacuum quantum state of the system. This definition leads to the usual relation

$$\langle 0 | g^2(\mathbf{x}, t) | 0 \rangle = \int \frac{dk}{k} \mathcal{P}_g(k). \quad (85)$$

6.5 Quantum fluctuations of a generic scalar field in a quasi de Sitter stage

So far, we have computed the time evolution and the spectrum of the quantum fluctuations of a generic scalar field χ supposing that the scale factor evolves like in a pure de Sitter expansion, $a(\tau) = -1/(H\tau)$. However, during inflation the Hubble rate is not exactly constant, but changes with time as $\dot{H} = -\epsilon H^2$ (quasi de Sitter expansion). In this subsection, we will solve for the perturbations in a quasi de Sitter expansion. Using the definition of the conformal time, one can show that the scale factor for small values of ϵ becomes

$$a(\tau) = -\frac{1}{H} \frac{1}{\tau(1-\epsilon)}.$$

Eq. (76) has now a squared mass term

$$M^2(\tau) = m_\chi^2 a^2 - \frac{a''}{a},$$

where

$$\begin{aligned} \frac{a''}{a} &= a^2 \left(\frac{\ddot{a}}{a} + H^2 \right) = a^2 (\dot{H} + 2H^2) \\ &= a^2 (2 - \epsilon) H^2 = \frac{(2 - \epsilon)}{\tau^2 (1 - \epsilon)^2} \\ &\simeq \frac{1}{\tau^2} (2 + 3\epsilon). \end{aligned} \quad (86)$$

Taking $m_\chi^2/H^2 = 3\eta_\chi$ and expanding for small values of ϵ and η we get Eq. (77) with

$$\nu_\chi \simeq \frac{3}{2} + \epsilon - \eta_\chi. \quad (87)$$

Armed with these results, we may compute the variance of the perturbations of the generic χ field

$$\begin{aligned}
\langle 0 | (\delta\chi(\mathbf{x}, t))^2 | 0 \rangle &= \int \frac{d^3k}{(2\pi)^3} |\delta\chi_{\mathbf{k}}|^2 \\
&= \int \frac{dk}{k} \frac{k^3}{2\pi^2} |\delta\chi_{\mathbf{k}}|^2 \\
&= \int \frac{dk}{k} \mathcal{P}_{\delta\chi}(k),
\end{aligned} \tag{88}$$

which defines the power spectrum of the fluctuations of the scalar field χ

$$\mathcal{P}_{\delta\chi}(k) \equiv \frac{k^3}{2\pi^2} |\delta\chi_{\mathbf{k}}|^2. \tag{89}$$

Since we have seen that fluctuations are (nearly) frozen in on superhorizon scales, a way of characterizing the perturbations is to compute the spectrum on scales larger than the horizon. For a massive scalar field, we obtain

$$\mathcal{P}_{\delta\chi}(k) = \left(\frac{H}{2\pi}\right)^2 \left(\frac{k}{aH}\right)^{3-2\nu_\chi}. \tag{90}$$

We may also define the *spectral index* $n_{\delta\chi}$ of the fluctuations as

$$n_{\delta\chi} - 1 = \frac{d\ln \mathcal{P}_{\delta\chi}}{d\ln k} = 3 - 2\nu_\chi = 2\eta_\chi - 2\epsilon.$$

The power spectrum of fluctuations of the scalar field χ is therefore *nearly flat*, that is is nearly independent from the wavelength $\lambda = \pi/k$: the amplitude of the fluctuation on superhorizon scales does not (almost) depend upon the time at which the fluctuations crosses the horizon and becomes frozen in. The small tilt of the power spectrum arises from the fact that the scalar field χ is massive and because during inflation the Hubble rate is not exactly constant, but nearly constant, where ‘nearly’ is quantified by the slow-roll parameters ϵ . Adopting the traditional terminology, we may say that the spectrum of perturbations is blue if $n_{\delta\chi} > 1$ (more power in the ultraviolet) and red if $n_{\delta\chi} < 1$ (more power in the infrared). The power spectrum of the perturbations of a generic scalar field χ generated during a period of slow roll inflation may be either blue or red. This depends upon the relative magnitude between η_χ and ϵ . For instance, in chaotic inflation with a quadric potential $V(\phi) = \frac{m_\phi^2 \phi^2}{2}$, one can easily compute

$$n_{\delta\chi} - 1 = 2\eta_\chi - 2\epsilon = \frac{2}{3H^2} (m_\chi^2 - m_\phi^2),$$

which tells us that the spectrum is blue (red) if $m_\chi^2 > m_\phi^2$ ($m_\chi^2 < m_\phi^2$).

Comment: We might have computed the spectral index of the spectrum $\mathcal{P}_{\delta\chi}(k)$ by first solving the equation for the perturbations of the field χ in a de Sitter stage, with $H = \text{constant}$ and therefore $\epsilon = 0$, and then taking into account the time-evolution of the Hubble rate introducing the subscript in H_k whose time variation is determined by Eq. (60). Correspondingly, H_k is the value of the Hubble rate when a given wavelength $\sim k^{-1}$ crosses the horizon (from that point on the fluctuations remains frozen in). The power spectrum in such an approach would read

$$\mathcal{P}_{\delta\chi}(k) = \left(\frac{H_k}{2\pi}\right)^2 \left(\frac{k}{aH}\right)^{3-2\nu_\chi} \quad (91)$$

with $3 - 2\nu_\chi \simeq \eta_\chi$. Using Eq. (60), one finds

$$n_{\delta\chi} - 1 = \frac{d\ln \mathcal{P}_{\delta\phi}}{d\ln k} = \frac{d\ln H_k^2}{d\ln k} + 3 - 2\nu_\chi = 2\eta_\chi - 2\epsilon$$

which reproduces our previous findings.

Comment: Since on superhorizon scales

$$\delta\chi_{\mathbf{k}} \simeq \frac{H}{\sqrt{2k^3}} \left(\frac{k}{aH}\right)^{\eta_\chi - \epsilon} \simeq \frac{H}{\sqrt{2k^3}} \left[1 + (\eta_\chi - \epsilon) \ln \left(\frac{k}{aH}\right)\right],$$

we discover that

$$|\delta\dot{\chi}_{\mathbf{k}}| \simeq |H(\eta_\chi - \epsilon) \delta\chi_{\mathbf{k}}| \ll |H \delta\chi_{\mathbf{k}}|, \quad (92)$$

that is on superhorizon scales the time variation of the perturbations can be safely neglected.

7 Quantum fluctuations during inflation

As we have mentioned in the previous section, the linear theory of the cosmological perturbations represent a cornerstone of modern cosmology and is used to describe the formation and evolution of structures in the universe as well as the anisotropies of the CMB. The seeds for these inhomogeneities were generated during inflation and stretched over astronomical scales because of the rapid superluminal expansion of the universe during the (quasi) de Sitter epoch.

In the previous section we have already seen that perturbations of a generic scalar field χ are generated during a (quasi) de Sitter expansion. The inflaton field is a scalar field and, as such, we conclude that inflaton fluctuations will be generated as well. However, the inflaton is special from the point of view of perturbations. The reason is very simple. By assumption, the inflaton field dominates the energy density of the universe during inflation. Any perturbation in the inflaton field means a perturbation of the stress energy-momentum tensor

$$\delta\phi \implies \delta T_{\mu\nu}.$$

A perturbation in the stress energy-momentum tensor implies, through Einstein's equations of motion, a perturbation of the metric

$$\delta T_{\mu\nu} \implies \left[\delta R_{\mu\nu} - \frac{1}{2} \delta (g_{\mu\nu} R) \right] = 8\pi G \delta T_{\mu\nu} \implies \delta g_{\mu\nu}.$$

On the other hand, a perturbation of the metric induces a backreaction on the evolution of the inflaton perturbation through the perturbed Klein-Gordon equation of the inflaton field

$$\delta g_{\mu\nu} \implies \delta \left(\partial_\mu \partial^\mu \phi + \frac{\partial V}{\partial \phi} \right) = 0 \implies \delta \phi.$$

This logic chain makes us conclude that the perturbations of the inflaton field and of the metric are tightly coupled to each other and have to be studied together

$$\delta \phi \iff \delta g_{\mu\nu}$$

As we will see shortly, this relation is stronger than one might thought because of the issue of gauge invariance.

Before launching ourselves into the problem of finding the evolution of the quantum perturbations of the inflaton field when they are coupled to gravity, let us give a heuristic explanation of why we expect that during inflation such fluctuations are indeed present.

If we take Eq. (50) and split the inflaton field as its classical value ϕ_0 plus the quantum fluctuation $\delta\phi$, $\phi(\mathbf{x}, t) = \phi_0(t) + \delta\phi(\mathbf{x}, t)$, the quantum perturbation $\delta\phi$ satisfies the equation of motion

$$\delta\ddot{\phi} + 3H \delta\dot{\phi} - \frac{\nabla^2 \delta\phi}{a^2} + V'' \delta\phi = 0. \quad (93)$$

Differentiating Eq. (55) wrt time and taking H constant (de Sitter expansion) we find

$$(\phi_0)''' + 3H\ddot{\phi}_0 + V'' \dot{\phi}_0 = 0. \quad (94)$$

Let us consider for simplicity the limit $\mathbf{k}^2/a^2 \ll 1$ and let us disregard the gradient term. Under this condition we see that $\dot{\phi}_0$ and $\delta\phi$ solve the same equation. The solutions have therefore to be related to each other by a constant of proportionality which depends upon time, that is

$$\delta\phi = -\dot{\phi}_0 \delta t(\mathbf{x}). \quad (95)$$

This tells us that $\phi(\mathbf{x}, t)$ will have the form

$$\phi(\mathbf{x}, t) = \phi_0(\mathbf{x}, t - \delta t(\mathbf{x})).$$

This equation indicates that the inflaton field does not acquire the same value at a given time t in all the space. On the contrary, when the inflaton field is rolling down its potential, it acquires different values from one spatial point $\mathbf{x}$ to the other. The inflaton field is not homogeneous and fluctuations are present. These fluctuations, in turn, will induce fluctuations in the metric.

7.1 The metric fluctuations

The mathematical tool to describe the linear evolution of the cosmological perturbations is obtained by perturbing at the first-order the FRW metric $g_{\mu\nu}^{(0)}$, see Eq. (1)

$$g_{\mu\nu} = g_{\mu\nu}^{(0)}(t) + g_{\mu\nu}(\mathbf{x}, t); \quad g_{\mu\nu} \ll g_{\mu\nu}^{(0)}. \quad (96)$$

The metric perturbations can be decomposed according to their spin with respect to a local rotation of the spatial coordinates on hypersurfaces of constant time. This leads to

- *scalar perturbations*
- *vector perturbations*
- *tensor perturbations*

Tensor perturbations or gravitational waves have spin 2 and are the “true” degrees of freedom of the gravitational fields in the sense that they can exist even in the vacuum. Vector perturbations are spin 1 modes arising from rotational velocity fields and are also called vorticity modes. Finally, scalar perturbations have spin 0.

Let us make a simple exercise to count how many scalar degrees of freedom are present. Take a spacetime of dimensions $D = n + 1$, of which n coordinates are spatial coordinates. The symmetric metric tensor $g_{\mu\nu}$ has $\frac{1}{2}(n + 2)(n + 1)$ degrees of freedom. We can perform $(n + 1)$ coordinate transformations in order to eliminate $(n + 1)$ degrees of freedom, this leaves us with $\frac{1}{2}n(n + 1)$ degrees of freedom. These $\frac{1}{2}n(n + 1)$ degrees of freedom contain scalar, vector and tensor modes. According to Helmholtz’s theorem we can always decompose a vector u_i ($i = 1, \dots, n$) as $u_i = \partial_i v + v_i$, where v is a scalar (usually called potential flow) which is curl-free, $v_{[i,j]} = 0$, and v_i is a real vector (usually called vorticity) which is divergence-free, $\nabla \cdot v = 0$. This means that the real vector (vorticity) modes are $(n - 1)$. Furthermore, a generic traceless tensor Π_{ij} can always be decomposed as $\Pi_{ij} = \Pi_{ij}^S + \Pi_{ij}^V + \Pi_{ij}^T$, where $\Pi_{ij}^S = \left(-\frac{k_i k_j}{k^2} + \frac{1}{3}\delta_{ij}\right) \Pi$, $\Pi_{ij}^V = (-i/2k)(k_i \Pi_j + k_j \Pi_i)$ ($k_i \Pi_i = 0$) and $k_i \Pi_{ij}^T = 0$. This means that the true symmetric, traceless and transverse tensor degrees of freedom are $\frac{1}{2}(n - 2)(n + 1)$.

The number of scalar degrees of freedom are therefore

$$\frac{1}{2}n(n + 1) - (n - 1) - \frac{1}{2}(n - 2)(n + 1) = 2,$$

while the degrees of freedom of true vector modes are $(n - 1)$ and the number of degrees of freedom of true tensor modes (gravitational waves) are $\frac{1}{2}(n - 2)(n + 1)$. In four dimensions $n = 3$, meaning that one expects 2 scalar degrees of freedom, 2 vector degrees of freedom and 2 tensor degrees of freedom. As we shall see, to the 2 scalar degrees of freedom from the metric, one has to add another one, the inflaton field perturbation $\delta\phi$. However, since Einstein’s equations will tell us that the two scalar degrees of freedom from the metric are equal during inflation, we expect a total number of scalar degrees of freedom equal to 2.

At the linear order, the scalar, vector and tensor perturbations evolve independently (they decouple) and it is therefore possible to analyze them separately. Vector perturbations are not excited during inflation because there are no rotational velocity fields during the

inflationary stage. We will analyze the generation of tensor modes (gravitational waves) in the following. For the time being we want to focus on the scalar degrees of freedom of the metric.

Considering only the scalar degrees of freedom of the perturbed metric, the most generic perturbed metric reads

$$g_{\mu\nu} = a^2 \begin{pmatrix} -1 - 2A & \partial_i B \\ \partial_i B & (1 - 2\psi)\delta_{ij} + D_{ij}E \end{pmatrix}, \quad (97)$$

while the line-element can be written as

$$ds^2 = a^2((-1 - 2A)d\tau^2 + 2\partial_i B d\tau dx^i + ((1 - 2\psi)\delta_{ij} + D_{ij}E) dx^i dx^j). \quad (98)$$

Here $D_{ij} = (\partial_i \partial_j - \frac{1}{3} \delta_{ij} \nabla^2)$.

We now want to determine the inverse $g^{\mu\nu}$ of the metric at the linear order

$$g^{\mu\alpha} g_{\alpha\nu} = \delta_\nu^\mu. \quad (99)$$

We have therefore to solve the equations

$$(g_{(0)}^{\mu\alpha} + g^{\mu\alpha}) (g_{\alpha\nu}^{(0)} + g_{\alpha\nu}) = \delta_\nu^\mu, \quad (100)$$

where $g_{(0)}^{\mu\alpha}$ is simply the unperturbed FRW metric (1). Since

$$g_{(0)}^{\mu\nu} = \frac{1}{a^2} \begin{pmatrix} -1 & 0 \\ 0 & \delta^{ij} \end{pmatrix}, \quad (101)$$

we can write in general

$$\begin{aligned} g^{00} &= \frac{1}{a^2} (-1 + X); \\ g^{0i} &= \frac{1}{a^2} \partial^i Y; \\ g^{ij} &= \frac{1}{a^2} ((1 + 2Z)\delta^{ij} + D^{ij}K). \end{aligned} \quad (102)$$

Plugging these expressions into Eq. (100) we find for $\mu = \nu = 0$

$$(-1 + X)(-1 - 2A) + \partial^i Y \partial_i B = 1. \quad (103)$$

Neglecting the terms $-2A \cdot X$ e $\partial^i Y \cdot \partial_i B$ because they are second-order in the perturbations, we find

$$1 - X + 2A = 1 \quad \Rightarrow \quad X = 2A. \quad (104)$$

Analogously, the components $\mu = 0, \nu = i$ of Eq. (100) give

$$(-1 + 2A)(\partial_i B) + \partial^j Y [(1 - 2\psi)\delta_{ji} + D_{ji}E] = 0. \quad (105)$$

At the first-order, we obtain

$$-\partial_i B + \partial_i Y = 0 \quad \Rightarrow \quad Y = B. \quad (106)$$

Finally, the components $\mu = i, \nu = j$ give

$$\partial^i B \partial_j B + \left((1 + 2Z)\delta^{ik} + D^{ik}K \right) ((1 - 2\psi)\delta_{kj} + D_{kj}E) = \delta_j^i. \quad (107)$$

Neglecting the second-order terms, we obtain

$$(1 - 2\psi + 2Z)\delta_j^i + D_j^i E + D_j^i K = \delta_j^i \Rightarrow Z = \psi; \quad K = -E. \quad (108)$$

The metric $g^{\mu\nu}$ finally reads

$$g^{\mu\nu} = \frac{1}{a^2} \begin{pmatrix} -1 + 2A & \partial^i B \\ \partial^i B & (1 + 2\psi)\delta^{ij} - D^{ij}E \end{pmatrix}. \quad (109)$$

7.2 Perturbed affine connections and Einstein's tensor

In this subsection we provide the reader with the perturbed affine connections and Einstein's tensor.

First, let us list the unperturbed affine connections

$$\Gamma_{00}^0 = \frac{a'}{a}; \quad \Gamma_{0j}^i = \frac{a'}{a} \delta_j^i; \quad \Gamma_{ij}^0 = \frac{a'}{a} \delta_{ij}; \quad (110)$$

$$\Gamma_{00}^i = \Gamma_{0i}^0 = \Gamma_{jk}^i = 0. \quad (111)$$

The expression for the affine connections in terms of the metric is

$$\Gamma_{\beta\gamma}^\alpha = \frac{1}{2} g^{\alpha\rho} \left(\frac{\partial g_{\rho\gamma}}{\partial x^\beta} + \frac{\partial g_{\beta\rho}}{\partial x^\gamma} - \frac{\partial g_{\beta\gamma}}{\partial x^\rho} \right) \quad (112)$$

which implies

$$\begin{aligned} \delta\Gamma_{\beta\gamma}^\alpha &= \frac{1}{2} \delta g^{\alpha\rho} \left(\frac{\partial g_{\rho\gamma}}{\partial x^\beta} + \frac{\partial g_{\beta\rho}}{\partial x^\gamma} - \frac{\partial g_{\beta\gamma}}{\partial x^\rho} \right) \\ &+ \frac{1}{2} g^{\alpha\rho} \left(\frac{\partial \delta g_{\rho\gamma}}{\partial x^\beta} + \frac{\partial \delta g_{\beta\rho}}{\partial x^\gamma} - \frac{\partial \delta g_{\beta\gamma}}{\partial x^\rho} \right), \end{aligned} \quad (113)$$

or in components

$$\delta\Gamma_{00}^0 = A'; \quad (114)$$

$$\delta\Gamma_{0i}^0 = \partial_i A + \frac{a'}{a} \partial_i B; \quad (115)$$

$$\delta\Gamma_{00}^i = \frac{a'}{a} \partial^i B + \partial^i B' + \partial^i A; \quad (116)$$

$$\begin{aligned} \delta\Gamma_{ij}^0 &= -2\frac{a'}{a}A\delta_{ij} - \partial_i\partial_j B - 2\frac{a'}{a}\psi\delta_{ij} \\ &\quad - \psi'\delta_{ij} - \frac{a'}{a}D_{ij}E + \frac{1}{2}D_{ij}E'; \end{aligned} \quad (117)$$

$$\delta\Gamma_{0j}^i = -\psi'\delta_{ij} + \frac{1}{2}D_{ij}E'; \quad (118)$$

$$\begin{aligned} \delta\Gamma_{jk}^i &= \partial_j\psi\delta_k^i - \partial_k\psi\delta_j^i + \partial^i\psi\delta_{jk} - \frac{a'}{a}\partial^i B\delta_{jk} \\ &\quad + \frac{1}{2}\partial_j D_k^i E + \frac{1}{2}\partial_k D_j^i E - \frac{1}{2}\partial^i D_{jk}E. \end{aligned} \quad (119)$$

We may now compute the Ricci scalar defines as

$$R_{\mu\nu} = \partial_\alpha \Gamma_{\mu\nu}^\alpha - \partial_\mu \Gamma_{\nu\alpha}^\alpha + \Gamma_{\sigma\alpha}^\alpha \Gamma_{\mu\nu}^\sigma - \Gamma_{\sigma\nu}^\alpha \Gamma_{\mu\alpha}^\sigma. \quad (120)$$

Its variation at the first-order reads

$$\begin{aligned} \delta R_{\mu\nu} &= \partial_\alpha \delta\Gamma_{\mu\nu}^\alpha - \partial_\mu \delta\Gamma_{\nu\alpha}^\alpha + \delta\Gamma_{\sigma\alpha}^\alpha \Gamma_{\mu\nu}^\sigma + \Gamma_{\sigma\alpha}^\alpha \delta\Gamma_{\mu\nu}^\sigma \\ &\quad - \delta\Gamma_{\sigma\nu}^\alpha \Gamma_{\mu\alpha}^\sigma - \Gamma_{\sigma\nu}^\alpha \delta\Gamma_{\mu\alpha}^\sigma. \end{aligned} \quad (121)$$

The background values are given by

$$R_{00} = -3\frac{a''}{a} + 3\left(\frac{a'}{a}\right)^2; \quad R_{0i} = 0; \quad (122)$$

$$R_{ij} = \left(\frac{a''}{a} + \left(\frac{a'}{a}\right)^2\right)\delta_{ij} \quad (123)$$

which give

$$\delta R_{00} = \frac{a'}{a}\partial_i\partial^i B + \partial_i\partial^i B' + \partial_i\partial^i A + 3\psi'' + 3\frac{a'}{a}\psi' + 3\frac{a'}{a}A'; \quad (124)$$

$$\delta R_{0i} = \frac{a''}{a}\partial_i B + \left(\frac{a'}{a}\right)^2\partial_i B + 2\partial_i\psi' + 2\frac{a'}{a}\partial_i A + \frac{1}{2}\partial_k D_i^k E'; \quad (125)$$

$$\begin{aligned} \delta R_{ij} &= \left(-\frac{a'}{a}A' - 5\frac{a'}{a}\psi' - 2\frac{a''}{a}A - 2\left(\frac{a'}{a}\right)^2 A\right. \\ &\quad \left.- 2\frac{a''}{a}\psi - 2\left(\frac{a'}{a}\right)^2\psi - \psi'' + \partial_k\partial^k\psi - \frac{a'}{a}\partial_k\partial^k B\right)\delta_{ij} \\ &\quad - \partial_i\partial_j B' + \frac{a'}{a}D_{ij}E' + \frac{a''}{a}D_{ij}E + \left(\frac{a'}{a}\right)^2 D_{ij}E \\ &\quad + \frac{1}{2}D_{ij}E'' + \partial_i\partial_j\psi - \partial_i\partial_j A - 2\frac{a'}{a}\partial_i\partial_j B \\ &\quad + \frac{1}{2}\partial_k\partial_i D_j^k E + \frac{1}{2}\partial_k\partial_j D_i^k E - \frac{1}{2}\partial_k\partial^k D_{ij}E; \end{aligned} \quad (126)$$

The perturbation of the scalar curvature

$$R = g^{\mu\alpha} R_{\alpha\mu}, \quad (127)$$

for which the first-order perturbation is

$$\delta R = \delta g^{\mu\alpha} R_{\alpha\mu} + g^{\mu\alpha} \delta R_{\alpha\mu}. \quad (128)$$

The background value is

$$R = \frac{6}{a^2} \frac{a''}{a} \quad (129)$$

while from Eq. (128) one finds

$$\begin{aligned} \delta R = & \frac{1}{a^2} \left(-6 \frac{a'}{a} \partial_i \partial^i B - 2 \partial_i \partial^i B' - 2 \partial_i \partial^i A - 6 \psi'' \right. \\ & \left. - 6 \frac{a'}{a} A' - 18 \frac{a'}{a} \psi' - 12 \frac{a''}{a} A + 4 \partial_i \partial^i \psi + \partial_k \partial^i D_i^k E \right). \end{aligned} \quad (130)$$

Finally, we may compute the perturbations of the Einstein tensor

$$G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R, \quad (131)$$

whose background components are

$$G_{00} = 3 \left(\frac{a'}{a} \right)^2; \quad G_{0i} = 0; \quad G_{ij} = \left(-2 \frac{a''}{a} + \left(\frac{a'}{a} \right)^2 \right) \delta_{ij}. \quad (132)$$

At first-order, one finds

$$\delta G_{\mu\nu} = \delta R_{\mu\nu} - \frac{1}{2} \delta g_{\mu\nu} R - \frac{1}{2} g_{\mu\nu} \delta R, \quad (133)$$

or in components

$$\delta G_{00} = -2 \frac{a'}{a} \partial_i \partial^i B - 6 \frac{a'}{a} \psi' + 2 \partial_i \partial^i \psi + \frac{1}{2} \partial_k \partial^i D_i^k E; \quad (134)$$

$$\delta G_{0i} = -2 \frac{a''}{a} \partial_i B + \left(\frac{a'}{a} \right)^2 \partial_i B + 2 \partial_i \psi' + \frac{1}{2} \partial_k D_i^k E' + 2 \frac{a'}{a} \partial_i A; \quad (135)$$

$$\begin{aligned} \delta G_{ij} = & \left(2 \frac{a'}{a} A' + 4 \frac{a'}{a} \psi' + 4 \frac{a''}{a} A - 2 \left(\frac{a'}{a} \right)^2 A \right. \\ & \left. + 4 \frac{a''}{a} \psi - 2 \left(\frac{a'}{a} \right)^2 \psi + 2 \psi'' - \partial_k \partial^k \psi \right) \delta_{ij} \end{aligned}$$

$$\begin{aligned}
& + 2 \frac{a'}{a} \partial_k \partial^k B + \partial_k \partial^k B' + \partial_k \partial^k A + \frac{1}{2} \partial_k \partial^m D_m^k E \Big) \delta_{ij} \\
& - \partial_i \partial_j B' + \partial_i \partial_j \psi - \partial_i \partial_j A + \frac{a'}{a} D_{ij} E' - 2 \frac{a''}{a} D_{ij} E \\
& + \left(\frac{a'}{a} \right)^2 D_{ij} E + \frac{1}{2} D_{ij} E'' + \frac{1}{2} \partial_k \partial_i D_j^k E \\
& + \frac{1}{2} \partial^k \partial_j D_{ik} E - \frac{1}{2} \partial_k \partial^k D_{ij} E - 2 \frac{a'}{a} \partial_i \partial_j B.
\end{aligned} \tag{136}$$

For convenience, we also give the expressions for the perturbations with one index up and one index down

$$\begin{aligned}
\delta G_\nu^\mu &= \delta(g^{\mu\alpha} G_{\alpha\nu}) \\
&= \delta g^{\mu\alpha} G_{\alpha\nu} + g^{\mu\alpha} \delta G_{\alpha\nu},
\end{aligned} \tag{137}$$

or in components

$$\delta G_0^0 = 6 \left(\frac{a'}{a} \right)^2 A + 6 \frac{a'}{a} \psi' + 2 \frac{a'}{a} \partial_i \partial^i B - 2 \partial_i \partial^i \psi - \frac{1}{2} \partial_k \partial^i D_i^k E. \tag{138}$$

$$\delta G_i^0 = -2 \frac{a'}{a} \partial_i A - 2 \partial_i \psi' - \frac{1}{2} \partial_k D_i^k E'. \tag{139}$$

$$\begin{aligned}
\delta G_j^i &= \left(2 \frac{a'}{a} A' + 4 \frac{a''}{a} A - 2 \left(\frac{a'}{a} \right)^2 A + \partial_i \partial^i A + 4 \frac{a'}{a} \psi' + 2 \psi'' \right. \\
&- \partial_i \partial^i \psi + 2 \frac{a'}{a} \partial_i \partial^i B + \partial_i \partial^i B' + \frac{1}{2} \partial_k \partial^m D_m^k E \Big) \delta_j^i \\
&- \partial^i \partial_j A + \partial^i \partial_j \psi - 2 \frac{a'}{a} \partial^i \partial_j B - \partial^i \partial_j B' + \frac{a'}{a} D_j^i E' + \frac{1}{2} D_j^i E'' \\
&+ \frac{1}{2} \partial_k \partial^i D_j^k E + \frac{1}{2} \partial_k \partial_j D^{ik} E - \frac{1}{2} \partial_k \partial^k D_j^i E.
\end{aligned} \tag{140}$$

7.3 Perturbed stress energy-momentum tensor

As we have seen previously, the perturbations of the metric are induced by the perturbations of the stress energy-momentum tensor of the inflaton field

$$T_{\mu\nu} = \partial_\mu \phi \partial_\nu \phi - g_{\mu\nu} \left(\frac{1}{2} g^{\alpha\beta} \partial_\alpha \phi \partial_\beta \phi - V(\phi) \right), \tag{141}$$

whose background values are

$$\begin{aligned}
T_{00} &= \frac{1}{2} \phi'^2 + V(\phi) a^2 \\
T_{0i} &= 0; \\
T_{ij} &= \left(\frac{1}{2} \phi'^2 - V(\phi) a^2 \right) \delta_{ij}.
\end{aligned} \tag{142}$$

The perturbed stress energy-momentum tensor reads

$$\begin{aligned}\delta T_{\mu\nu} &= \partial_\mu \delta\phi \partial_\nu \phi + \partial_\mu \phi \partial_\nu \delta\phi - \delta g_{\mu\nu} \left(\frac{1}{2} g^{\alpha\beta} \partial_\alpha \phi \partial_\beta \phi + V(\phi) \right) \\ &- g_{\mu\nu} \left(\frac{1}{2} \delta g^{\alpha\beta} \partial_\alpha \phi \partial_\beta \phi + g^{\alpha\beta} \partial_\alpha \delta\phi \partial_\beta \phi + \frac{\partial V}{\partial \phi} \delta\phi + \frac{\partial V}{\partial \phi} \delta\phi \right).\end{aligned}\quad (143)$$

In components we have

$$\delta T_{00} = \delta\phi' \phi' + 2A V(\phi) a^2 + a^2 \frac{\partial V}{\partial \phi} \delta\phi;\quad (144)$$

$$\delta T_{0i} = \partial_i \delta\phi \phi' + \frac{1}{2} \partial_i B \phi'^2 - \partial_i B V(\phi) a^2;\quad (145)$$

$$\begin{aligned}\delta T_{ij} &= \left(\delta\phi' \phi' - A \phi'^2 - a^2 \frac{\partial V}{\partial \phi} \delta^{(1)}\phi - \psi \phi'^2 + 2\psi V(\phi) a^2 \right) \delta_{ij} \\ &+ \frac{1}{2} D_{ij} E \phi'^2 - D_{ij} E V(\phi) a^2.\end{aligned}\quad (146)$$

For convenience, we list the mixed components

$$\begin{aligned}\delta T_\nu^\mu &= \delta(g^{\mu\alpha} T_{\alpha\nu}) \\ &= \delta g^{\mu\alpha} T_{\alpha\nu} + g^{\mu\alpha} \delta T_{\alpha\nu}\end{aligned}\quad (147)$$

or

$$\begin{aligned}\delta T_0^0 &= A \phi'^2 - \delta\phi' \phi' - \delta\phi \frac{\partial V}{\partial \phi} a^2; \\ \delta T_0^i &= \partial^i B \phi'^2 + \partial^i \delta\phi \phi'; \\ \delta T_i^0 &= -\partial^i \delta\phi \phi'; \\ \delta T_j^i &= \left(-A \phi'^2 + \delta\phi' \phi' - \delta\phi \frac{\partial V}{\partial \phi} a^2 \right) \delta_j^i.\end{aligned}\quad (148)$$

7.4 Perturbed Klein-Gordon equation

The inflaton equation of motion is the Klein-Gordon equation of a scalar field under the action of its potential $V(\phi)$. The equation to perturb is therefore

$$\begin{aligned}\partial^\mu \partial_\mu \phi &= \frac{\partial V}{\partial \phi}; \\ \partial_\mu \partial^\mu \phi &= \frac{1}{\sqrt{-g}} \partial_\nu \left(\sqrt{-g} g^{\mu\nu} \partial_\nu \phi \right);\end{aligned}\quad (149)$$

which at the zero-th order gives the inflaton equation of motion

$$\phi'' + 2 \frac{a'}{a} \phi' = - \frac{\partial V}{\partial \phi} a^2.\quad (150)$$

The variation of Eq. (149) is the sum of four different contributions corresponding to the variations of $\frac{1}{\sqrt{-g}}$, $\sqrt{-g}$, $g^{\mu\nu}$ and ϕ . For the variation of g we have

$$\delta g = g g^{\mu\nu} \delta g_{\nu\mu} = dg = g^{\mu\nu} dg_{\mu\nu} \quad (151)$$

which give at the linear order

$$\begin{aligned} \delta\sqrt{-g} &= -\frac{\delta g}{2\sqrt{-g}}; \\ \delta\frac{1}{\sqrt{-g}} &= \frac{\delta\sqrt{-g}}{g}. \end{aligned} \quad (152)$$

Plugging these results into the expression for the variation of Eq. (150)

$$\begin{aligned} \delta\partial_\mu\partial^\mu\phi &= -\delta\phi'' - 2\frac{a'}{a}\delta\phi' + \partial_i\partial^i\delta\phi + 2A\phi'' + 4\frac{a'}{a}A\phi' + A'\phi' \\ &+ 3\psi'\phi' + \partial_i\partial^i B\phi' \\ &= \delta\phi\frac{\partial^2 V}{\partial\phi^2}a^2. \end{aligned} \quad (153)$$

Using Eq. (150) to write

$$2A\phi'' + 4\frac{a'}{a}\phi' = 2A\frac{\partial V}{\partial\phi}, \quad (154)$$

Eq. (153) becomes

$$\begin{aligned} \delta\phi'' + 2\frac{a'}{a}\delta\phi' &- \partial_i\partial^i\delta\phi - A'\phi' - 3\psi'\phi' - \partial_i\partial^i B\phi' \\ &= -\delta\phi\frac{\partial^2 V}{\partial\phi^2}a^2 - 2A\frac{\partial V}{\partial\phi}. \end{aligned} \quad (155)$$

After having computed the perturbations at the linear order of the Einstein's tensor and of the stress energy-momentum tensor, we are ready to solve the perturbed Einstein's equations in order to quantify the inflaton and the metric fluctuations. We pause, however, for a moment in order to deal with the problem of gauge invariance.

7.5 The issue of gauge invariance

When studying the cosmological density perturbations, what we are interested in is following the evolution of a spacetime which is neither homogeneous nor isotropic. This is done by following the evolution of the differences between the actual spacetime and a well understood reference spacetime. So we will consider small perturbations away from the homogeneous, isotropic spacetime (see Fig. 9). The reference system in our case is the spatially flat Friedmann–Robertson–Walker (FRW) spacetime, with line element $ds^2 = a^2(\tau)\{d\tau^2 - \delta_{ij}dx^i dx^j\}$. Now, the key issue is that general relativity is a gauge theory where the gauge transformations are the generic coordinate transformations from a local reference frame to another.

When we compute the perturbation of a given quantity, this is defined to be the difference between the value that this quantity assumes on the real physical spacetime and the value it assumes on the unperturbed background. Nonetheless, to perform a comparison between these two values, it is necessary to compute the at the same spacetime point. Since the two values “live” on two different geometries, it is necessary to specify a map which allows to link univocally the same point on the two different spacetimes. This correspondance is called a gauge choice and changing the map means performing a gauge transformation.

Fixing a gauge in general relativity implies choosing a coordinate system. A choice of coordinates defines a *threading* of spacetime into lines (corresponding to fixed spatial coordinates $\mathbf{x}$) and a *slicing* into hypersurfaces (corresponding to fixed time τ). A choice of coordinates is called a *gauge* and there is no unique preferred gauge

GAUGE CHOICE $\iff$ SLICING AND THREADING

From a more formal point of view, operating an infinitesimal gauge tranformation on the coordinates

$$\widetilde{x}^\mu = x^\mu + \delta x^\mu \quad (156)$$

implies on a generic quantity Q a transformation on its perturbation

$$\delta \widetilde{Q} = \delta Q + \mathcal{L}_{\delta x} Q_0 \quad (157)$$

where Q_0 is the value assumed by the quantity Q on the background and $\mathcal{L}_{\delta x}$ is the Lie-derivative of Q along the vector δx^μ .

Decomposing in the usual manner the vector δx^μ

$$\begin{aligned} \delta x^0 &= \xi^0(x^\mu); \\ \delta x^i &= \partial^i \beta(x^\mu) + v^i(x^\mu); \quad \partial_i v^i = 0, \end{aligned} \quad (158)$$

we can easily deduce the transformation law of a scalar quantity f (like the inflaton scalar field ϕ and energy density ρ). Instead of applying the formal definition (157), we find the transformation law in an alternative (and more pedagogical) way. We first write $\delta f(x) = \widetilde{f}(x) - f_0(x)$, where $f_0(x)$ is the background value. Under a gauge transformation we have $\delta \widetilde{f}(\widetilde{x}^\mu) = \widetilde{f}(\widetilde{x}^\mu) - \widetilde{f}_0(\widetilde{x}^\mu)$. Since f is a scalar we can write $f(\widetilde{x}^\mu) = f(x^\mu)$ (the value of the scalar function in a given physical point is the same in all the coordinate system). On the other side, on the unperturbed background hypersurface $\widetilde{f}_0 = f_0$. We have therefore

$$\begin{aligned} \delta \widetilde{f}(\widetilde{x}^\mu) &= \widetilde{f}(\widetilde{x}^\mu) - \widetilde{f}_0(\widetilde{x}^\mu) \\ &= f(x^\mu) - f_0(\widetilde{x}^\mu) \\ &= f(\widetilde{x}^\mu) - f_0(\widetilde{x}^\mu) \\ &= f(\widetilde{x}^\mu) - \delta x^\mu \frac{\partial f}{\partial x^\mu}(\widetilde{x}) - f_0(\widetilde{x}^\mu), \end{aligned} \quad (159)$$

from which we finally deduce, being $f_0 = f_0(x^0)$,

$$\widetilde{\delta f} = \delta f - f' \xi^0$$

For the spin zero perturbations of the metric, we can proceed analogously. We use the following trick. Upon a coordinate transformation $x^\mu \rightarrow \widetilde{x}^\mu = x^\mu + \delta x^\mu$, the line element is left invariant, $ds^2 = \widetilde{ds}^2$. This implies, for instance, that $a^2(\widetilde{x}^0) (1 + \widetilde{A}) (d\widetilde{x}^0)^2 = a^2(x^0) (1 + A) (dx^0)^2$. Since $a^2(\widetilde{x}^0) \simeq a^2(x^0) + 2a a' \xi^0$ and $d\widetilde{x}^0 = (1 + \xi^{0'}) dx^0 + \frac{\partial x^0}{\partial x^i} dx^i$, we obtain $1 + 2A = 1 + 2\widetilde{A} + 2\mathcal{H}\xi^0 + 2\xi^{0'}$. A similar procedure leads to the following transformation laws

$$\begin{aligned} \widetilde{A} &= A - \xi^{0'} - \frac{a'}{a} \xi^0; \\ \widetilde{B} &= B + \xi^0 + \beta' \\ \widetilde{\psi} &= \psi - \frac{1}{3} \nabla^2 \beta + \frac{a'}{a} \xi^0; \\ \widetilde{E} &= E + 2\beta. \end{aligned}$$

The gauge problem stems from the fact that a change of the map (a change of the coordinate system) implies the variation of the perturbation of a given quantity which may therefore assume different values (all of them on a equal footing!) according to the gauge choice. To eliminate this ambiguity, one has therefore a double choice:

- Identify those combinations representing gauge invariant quantities;
- choose a given gauge and perform the calculations in that gauge.

Both options have advantages and drawbacks. Choosing a gauge may render the computation technically simpler with the danger, however, of including gauge artifacts, *i.e.* gauge freedoms which are not physical. Performing a gauge-invariant computation may be technically more involved, but has the advantage of treating only physical quantities.

Let us first indicate some gauge-invariant quantities which have been introduced first in Ref. [7]. They are the so-called gauge invariant potentials or Bardeen's potentials

$$\Phi = -A + \frac{1}{a} \left[\left(-B + \frac{E'}{2} \right) a \right]', \quad (160)$$

$$\Psi = -\psi - \frac{1}{6} \nabla^2 E + \frac{a'}{a} \left(B - \frac{E'}{2} \right). \quad (161)$$

Analogously, one can define a gauge invariant quantity for the perturbation of the inflaton field. Since ϕ is a scalar field $\widetilde{\delta\phi} = (\delta\phi - \phi' \xi^0)$ and therefore

$$\delta\phi^{(\text{GI})} = -\delta\phi + \phi' \left(\frac{E'}{2} - B \right)$$

is gauge-invariant.

Analogously, one can define a gauge-invariant energy-density perturbation

$$\delta\rho^{(\text{GI})} = -\delta\rho + \rho' \left(\frac{E'}{2} - B \right)$$

We now want to pause to introduce in details some gauge-invariant quantities which play a major role in the computation of the density perturbations. In the following we will be interested only in the coordinate transformations on constant time hypersurfaces and therefore gauge invariance will be equivalent to independent of the slicing.

7.6 The comoving curvature perturbation

The intrinsic spatial curvature on hypersurfaces on constant conformal time τ and for a flat universe is given by

$$^{(3)}R = \frac{4}{a^2} \nabla^2 \psi.$$

The quantity ψ is usually referred to as the *curvature perturbation*. We have seen, however, that the curvature potential ψ is *not* gauge invariant, but is defined only on a given slicing. Under a transformation on constant time hypersurfaces $t \rightarrow t + \delta\tau$ (change of the slicing)

$$\psi \rightarrow \psi + \mathcal{H} \delta\tau.$$

We now consider the *comoving slicing* which is defined to be the slicing orthogonal to the worldlines of comoving observers. The latter are free-falling and the expansion defined by them is isotropic. In practice, what this means is that there is no flux of energy measured by these observers, that is $T_{0i} = 0$. During inflation this means that these observers measure $\delta\phi_{\text{com}} = 0$ since T_{0i} goes like $\partial_i \delta\phi(\mathbf{x}, \tau) \phi'(\tau)$.

Since $\delta\phi \rightarrow \delta\phi - \phi' \delta\tau$ for a transformation on constant time hypersurfaces, this means that

$$\delta\phi \rightarrow \delta\phi_{\text{com}} = \delta\phi - \phi' \delta\tau = 0 \implies \delta\tau = \frac{\delta\phi}{\phi'},$$

that is $\delta\tau = \frac{\delta\phi}{\phi'}$ is the time-displacement needed to go from a generic slicing with generic $\delta\phi$ to the comoving slicing where $\delta\phi_{\text{com}} = 0$. At the same time the curvature perturbation ψ transforms into

$$\psi \rightarrow \psi_{\text{com}} = \psi + \mathcal{H} \delta\tau = \psi + \mathcal{H} \frac{\delta\phi}{\dot{\phi}}.$$

The quantity

$$\mathcal{R} = \psi + \mathcal{H} \frac{\delta\phi}{\dot{\phi}} = \psi + H \frac{\delta\phi}{\dot{\phi}}$$

is the *comoving curvature perturbation*. This quantity is gauge invariant by construction and is related to the gauge-dependent curvature perturbation ψ on a generic slicing to the inflaton perturbation $\delta\phi$ in that gauge. By construction, the meaning of $\mathcal{R}$ is that it represents the gravitational potential on comoving hypersurfaces where $\delta\phi = 0$

$$\mathcal{R} = \psi|_{\delta\phi=0}.$$

7.7 The curvature perturbation on spatial slices of uniform energy density

We now consider the *slicing of uniform energy density* which is defined to be the the slicing where there is no perturbation in the energy density, $\delta\rho = 0$.

Since $\delta\rho \rightarrow \delta\rho - \rho' \delta\tau$ for a transformation on constant time hypersurfaces, this means that

$$\delta\rho \rightarrow \delta\rho_{\text{unif}} = \delta\rho - \rho' \delta\tau = 0 \implies \delta\tau = \frac{\delta\rho}{\rho'},$$

that is $\delta\tau = \frac{\delta\rho}{\rho'}$ is the time-displacement needed to go from a generic slicing with generic $\delta\rho$ to the slicing of uniform energy density where $\delta\rho_{\text{unif}} = 0$. At the same time the curvature perturbation ψ transforms into

$$\psi \rightarrow \psi_{\text{unif}} = \psi + \mathcal{H} \delta\tau = \psi + \mathcal{H} \frac{\delta\rho}{\rho'}.$$

The quantity

$$\zeta = \psi + \mathcal{H} \frac{\delta\rho}{\rho'} = \psi + H \frac{\delta\rho}{\dot{\rho}}$$

is the *curvature perturbation on slices of uniform energy density*. This quantity is gauge invariant by construction and is related to the gauge-dependent curvature perturbation ψ on a generic slicing and to the energy density perturbation $\delta\rho$ in that gauge. By construction,

the meaning of ζ is that it represents the gravitational potential on slices of uniform energy density

$$\zeta = \psi|_{\delta\rho=0}.$$

Notice that, using the energy-conservation equation $\rho' + 3\mathcal{H}(\rho + p) = 0$, the curvature perturbation on slices of uniform energy density can be also written as

$$\zeta = \psi - \frac{\delta\rho}{3(\rho + p)}.$$

During inflation $\rho + p = \dot{\phi}^2$. Furthermore, on superhorizon scales from what we have learned in the previous section (and will be rigously shown in the following) the inflaton fluctuation $\delta\phi$ is frozen in and $\delta\dot{\phi} = (\text{slow roll parameters}) \times H \delta\phi$. This implies that $\delta\rho = \dot{\phi}\delta\dot{\phi} + V'\delta\phi \simeq V'\delta\phi \simeq -3H\dot{\phi}$, leading to

$$\zeta \simeq \psi + \frac{3H\dot{\phi}}{3\dot{\phi}^2} = \psi + H\frac{\delta\phi}{\dot{\phi}}\mathcal{R} \quad (\text{ON SUPERHORIZON SCALES})$$

The comoving curvature perturbation and the curvature perturbation on uniform energy density slices are equal on superhorizon scales.

7.8 Scalar field perturbations in the spatially flat gauge

We now consider the *spatially flat gauge* which is defined to be the the slicing where there is no curvature $\psi_{\text{flat}} = 0$.

Since $\psi \rightarrow \psi + \mathcal{H}\delta\tau$ for a transformation on constant time hypersurfaces, this means that

$$\psi \rightarrow \psi_{\text{flat}} = \psi + \mathcal{H}\delta\tau = 0 \implies \delta\tau = -\frac{\psi}{\mathcal{H}},$$

that is $\delta\tau = -\psi/\mathcal{H}$ is the time-displacement needed to go from a generic slicing with generic ψ to the spatially flat gauge where $\psi_{\text{flat}} = 0$. At the same time the fluctuation of the inflaton field transforms a

$$\delta\phi \rightarrow \delta\phi - \phi' \delta\tau = \delta\phi + \frac{\phi'}{\mathcal{H}} \psi.$$

The quantity

$$Q = \delta\phi + \frac{\phi'}{\mathcal{H}} \psi = \delta\phi + \frac{\dot{\phi}}{H} \psi \equiv \frac{\dot{\phi}}{H} \mathcal{R}$$

is the inflaton perturbation on spatially flat gauges. This quantity is gauge invariant by construction and is related to the inflaton perturbation $\delta\phi$ on a generic slicing and to to

the curvature perturbation ψ in that gauge. By construction, the meaning of Q is that it represents the inflaton potential on spatially flat slices

$$Q = \delta\phi|_{\delta\psi=0}.$$

This quantity is often referred to as the Sasaki or Mukhanov variable [62, 52].

Notice that $\delta\phi = -\phi'\delta\tau = -\dot{\phi}\delta t$ on flat slices, where δt is the time displacement going from flat to comoving slices. This relation makes somehow rigorous the expression (95). Analogously, going from flat to comoving slices one has $\mathcal{R} = H\delta t$.

7.9 Adiabatic and isocurvature perturbations

Arbitrary cosmological perturbations can be decomposed into:

- *adiabatic or curvature perturbations* which perturb the solution along the same trajectory in phase-space as the as the background solution. The perturbations in any scalar quantity X can be described by a unique perturbation in expansion with respect to the background

$$H\delta t = H\frac{\delta X}{\dot{X}} \quad \text{FOR EVERY } X.$$

In particular, this holds for the energy density and the pressure

$$\frac{\delta\rho}{\dot{\rho}} = \frac{\delta p}{\dot{p}}$$

which implies that $p = p(\rho)$. This explains why they are called adiabatic. They are called curvature perturbations because a given time displacement δt causes the same relative change $\delta X/\dot{X}$ for all quantities. In other words the perturbations is democratically shared by all components of the universe.

- *isocurvature perturbations* which perturb the solution off the background solution

$$\frac{\delta X}{\dot{X}} \neq \frac{\delta Y}{\dot{Y}} \quad \text{FOR SOME } X \text{ AND } Y.$$

One way of specifying a generic isocurvature perturbation δX is to give its value on uniform-density slices, related to its value on a different slicing by the gauge-invariant definition

$$H\left.\frac{\delta X}{\dot{X}}\right|_{\delta\rho=0} = H\left(\frac{\delta X}{\dot{X}} - \frac{\delta\rho}{\dot{\rho}}\right).$$

For a set of fluids with energy density ρ_i , the isocurvature perturbations are conventionally defined by the gauge invariant quantities

$$S_{ij} = 3H \left(\frac{\delta\rho_i}{\dot{\rho}_i} - \frac{\delta\rho_j}{\dot{\rho}_j} \right) = 3(\zeta_i - \zeta_j).$$

One simple example of isocurvature perturbations is the baryon-to-photon ratio $S = \delta(n_B/n_\gamma) = (\delta n_B/n_B) - (\delta n_\gamma/n_\gamma)$.

1. *Comment:*

From the definitions above, it follows that the cosmological perturbations generated during inflation are of the adiabatic type *if* the inflaton field is the only field driving inflation. However, if inflation is driven by more than one field, isocurvature perturbations are expected to be generated (and they might even be cross-correlated to the adiabatic ones [9, 10, 12]). In the following, however, we will keep focussing – as done so far – on the one-single field case, that is we will be dealing only with adiabatic/curvature perturbations.

2. *Comment:* The perturbations generated during inflation are *gaussian*, *i.e.* the two-point correlation functions (like the power spectrum) suffice to define all the higher-order even correlation functions, while the odd correlation functions (such as the three-point correlation function) vanish. This conclusion is drawn by the very same fact that cosmological perturbations are studied *linearizing* Einstein’s and Klein-Gordon equations. This turns out to be a good approximation because we know that the inflaton potential needs to be very flat in order to drive inflation and the interaction terms in the inflaton potential might be present, but they are small. Non-gaussian features are therefore suppressed since the non-linearities of the inflaton potential are suppressed too. The same argument applies to the metric perturbations; non-linearities appear only at the second-order in deviations from the homogeneous background solution and are therefore small. This expectation has been recently confirmed by the first computation of the cosmological perturbations generated during inflation up to second-order in deviations from the homogeneous background solution which fully account for the inflaton self-interactions as well as for the second-order fluctuations of the background metric [2].

7.10 The next steps

After all these technicalities, it is useful to rest for a moment and to go back to physics. Up to now we have learned that during inflation quantum fluctuations of the inflaton field are generated and their wavelengths are stretched on large scales by the rapid expansion of the universe. We have also seen that the quantum fluctuations of the inflaton field are in fact impossible to disentangle from the metric perturbations. This happens not only because they are tightly coupled to each other through Einstein’s equations, but also because of the issue of gauge invariance. Take, for instance, the gauge invariant quantity $Q = \delta\phi + \frac{\phi'}{\mathcal{H}} \psi$. We can always go to a gauge where the fluctuation is entirely in the curvature potential ψ , $Q = \frac{\phi'}{\mathcal{H}} \psi$, or entirely in the inflaton field, $Q = \delta\phi$. However, as we have stressed at the end of the previous section, once ripples in the curvature are frozen in on superhorizon scales during inflation, it is in fact gravity that acts as a messenger communicating to baryons and photons

the small seeds of perturbations once a given scale reenters the horizon after inflation. This happens thanks to Newtonian physics; a small perturbation in the gravitational potential ψ induces a small perturbation of the energy density ρ through Poisson's equation $\nabla^2\psi = 4\pi G\delta\rho$. Similarly, if perturbations are adiabatic/curvature perturbations and, as such, treat democratically all the components, a ripple in the curvature is communicated to photons as well, giving rise to a nonvanishing $\delta T/T$.

These considerations make it clear that the next steps of these lectures will be

- Compute the curvature perturbation generated during inflation on superhorizon scales. As we have seen we can either compute the comoving curvature perturbation $\mathcal{R}$ or the curvature on uniform energy density hypersurfaces ζ . They will tell us about the fluctuations of the gravitational potential.
- See how the fluctuations of the gravitational potential are transmitted to baryons and photons.

We now intend to address the first point. As stressed previously, we are free to follow two alternative roads: either pick up a gauge and compute the gauge-invariant curvature in that gauge or perform a gauge-invariant calculation. We take both options.

7.11 Computation of the curvature perturbation using the longitudinal gauge

The longitudinal (or conformal newtonian) gauge is a convenient gauge to compute the cosmological perturbations. It is defined by performing a coordinate transformation such that $B = E = 0$. This leaves behind two degrees of freedom in the scalar perturbations, A and ψ . As we have previously seen in subsection 7.1, these two degrees of freedom fully account for the scalar perturbations in the metric.

First of all, we take the non-diagonal part ($i \neq j$) of Eq. (140). Since the stress energy-momentum tensor does not have any non-diagonal component (no stress), we have

$$\partial_i \partial_j (\psi - A) = 0 \implies \psi = A$$

and we can now work only with one variable, let it be ψ .

Eq. (139) gives (in cosmic time)

$$\dot{\psi} + H\psi = 4\pi G\dot{\phi}\delta\phi = \epsilon H \frac{\delta\phi}{\dot{\phi}}, \quad (162)$$

while Eq. (138) and the diagonal part of (140) ($i = j$) give respectively

$$-3H(\dot{\psi} + H\psi) + \frac{\nabla^2\psi}{a^2} = 4\pi G(\dot{\phi}\delta\dot{\phi} - \dot{\phi}^2\psi + V'\delta\phi), \quad (163)$$

$$-\left(2\frac{\ddot{a}}{a} + \left(\frac{\dot{a}}{a}\right)^2\right)\psi - 3H\dot{\psi} - \ddot{\psi} = -(\dot{\phi}\delta\dot{\phi} - \dot{\phi}^2\psi - V'\delta\phi), \quad (164)$$

If we now use the fact that $\dot{H} = 4\pi G\dot{\phi}^2$, sum Eqs. (163) and 164) and use the background Klein-Gordon equation to eliminate V' , we arrive at the equation for the gravitational potential

$$\ddot{\psi}_{\mathbf{k}} + \left(H - 2\frac{\ddot{\phi}}{\dot{\phi}}\right)\dot{\psi}_{\mathbf{k}} + 2\left(\dot{H} - H\frac{\ddot{\phi}}{\dot{\phi}}\right)\psi_{\mathbf{k}} + \frac{k^2}{a^2}\psi_{\mathbf{k}} = 0 \quad (165)$$

We may rewrite it in conformal time

$$\psi_{\mathbf{k}}'' + 2\left(\mathcal{H} - \frac{\phi''}{\phi'}\right)\psi_{\mathbf{k}}' + 2\left(\mathcal{H}' - \mathcal{H}\frac{\phi''}{\phi'}\right)\psi_{\mathbf{k}} + k^2\psi_{\mathbf{k}} = 0 \quad (166)$$

and in terms of the slow-roll parameters ϵ and η

$$\psi_{\mathbf{k}}'' + 2\mathcal{H}(\eta - \epsilon)\psi_{\mathbf{k}}' + 2\mathcal{H}^2(\eta - 2\epsilon)\psi_{\mathbf{k}} + k^2\psi_{\mathbf{k}} = 0. \quad (167)$$

Using the same logic leading to Eq. (92), from Eq. (165) we can infer that on superhorizon scales the gravitational potential ψ is nearly constant (up to a mild logarithmic time-dependence proportional to slow-roll parameters), that is $\dot{\psi}_{\mathbf{k}} \sim (\text{slow-roll parameters}) \times \psi_{\mathbf{k}}$. This is hardly surprising, we know that fluctuations are frozen in on superhorizon scales.

Using Eq. (162), we can therefore relate the fluctuation of the gravitational potential ψ to the fluctuation of the inflaton field $\delta\phi$ on superhorizon scales

$$\psi_{\mathbf{k}} \simeq \epsilon H \frac{\delta\phi_{\mathbf{k}}}{\dot{\phi}} \quad (\text{ON SUPERHORIZON SCALES}) \quad (168)$$

This gives us the chance to compute the gauge-invariant comoving curvature perturbation $\mathcal{R}_{\mathbf{k}}$

$$\mathcal{R}_{\mathbf{k}} = \psi_{\mathbf{k}} + H \frac{\delta\phi_{\mathbf{k}}}{\dot{\phi}} = (1 + \epsilon) \frac{\delta\phi_{\mathbf{k}}}{\dot{\phi}} \simeq \frac{\delta\phi_{\mathbf{k}}}{\dot{\phi}}. \quad (169)$$

The power spectrum of the the comoving curvature perturbation $\mathcal{R}_{\mathbf{k}}$ then reads on superhorizon scales

$$\mathcal{P}_{\mathcal{R}} = \frac{k^3}{2\pi^2} \frac{H^2}{\dot{\phi}^2} |\delta\phi_{\mathbf{k}}|^2 = \frac{k^3}{4m_{\text{Pl}}^2 \epsilon \pi^2} |\delta\phi_{\mathbf{k}}|^2.$$

What is left to evaluate is the time evolution of $\delta\phi_{\mathbf{k}}$. To do so, we consider the perturbed Klein-Gordon equation (155) in the longitudinal gauge (in cosmic time)

$$\delta\ddot{\phi}_{\mathbf{k}} + 3H\delta\dot{\phi}_{\mathbf{k}} + \frac{k^2}{a^2}\delta\phi_{\mathbf{k}} + V''\delta\phi_{\mathbf{k}} = -2\psi_{\mathbf{k}}V' + 4\dot{\psi}_{\mathbf{k}}\dot{\phi}.$$

Since on superhorizon scales $|4\dot{\psi}_{\mathbf{k}}\dot{\phi}| \ll |\psi_{\mathbf{k}}V'|$, using Eq. (168) and the relation $V' \simeq -3H\dot{\phi}$, we can rewrite the perturbed Klein-Gordon equation on superhorizon scales as

$$\delta\ddot{\phi}_{\mathbf{k}} + 3H\delta\dot{\phi}_{\mathbf{k}} + (V'' + 6\epsilon H^2)\delta\phi_{\mathbf{k}} = 0.$$

We now introduce as usual the field $\delta\chi_{\mathbf{k}} = \delta\phi_{\mathbf{k}}/a$ and go to conformal time τ . The perturbed Klein-Gordon equation on superhorizon scales becomes, using Eq. (86),

$$\begin{aligned}\delta\chi_{\mathbf{k}}'' &- \frac{1}{\tau^2} \left(\nu^2 - \frac{1}{4} \right) \delta\chi_{\mathbf{k}} = 0, \\ \nu^2 &= \frac{9}{4} + 9\epsilon - 3\eta.\end{aligned}\tag{170}$$

Using what we have learned in the previous section, we conclude that

$$|\delta\phi_{\mathbf{k}}| \simeq \frac{H}{\sqrt{2k^3}} \left(\frac{k}{aH} \right)^{\frac{3}{2}-\nu} \quad (\text{ON SUPERHORIZON SCALES})$$

which justifies our initial assumption that both the inflaton perturbation and the gravitational potential are nearly constant on superhorizon scale.

We may now compute the power spectrum of the comoving curvature perturbation on superhorizon scales

$$\mathcal{P}_{\mathcal{R}}(k) = \frac{1}{2m_{\text{Pl}}^2\epsilon} \left(\frac{H}{2\pi} \right)^2 \left(\frac{k}{aH} \right)^{n_{\mathcal{R}}-1} \equiv A_{\mathcal{R}}^2 \left(\frac{k}{aH} \right)^{n_{\mathcal{R}}-1}$$

where we have defined the *spectral index* $n_{\mathcal{R}}$ of the comoving curvature perturbation as

$$n_{\mathcal{R}} - 1 = \frac{d \ln \mathcal{P}_{\mathcal{R}}}{d \ln k} = 3 - 2\nu = 2\eta - 6\epsilon.$$

We conclude that inflation is responsible for the generation of adiabatic/curvature perturbations with an almost scale-independent spectrum.

From the curvature perturbation we can easily deduce the behaviour of the gravitational potential $\psi_{\mathbf{k}}$ from Eq. (162). The latter is solved by

$$\psi_{\mathbf{k}} = \frac{A(k)}{a} + \frac{4\pi G}{a} \int^t dt' a(t') \dot{\phi}(t') \delta\phi_{\mathbf{k}}(t') \simeq \frac{A(k)}{a} + \epsilon \mathcal{R}_{\mathbf{k}}.$$

We find that during inflation and on superhorizon scales the gravitational potential is the sum of a decreasing function plus a nearly constant in time piece proportional to the curvature perturbation. Notice in particular that in an exact de Sitter stage, that is $\epsilon = 0$, the gravitational potential is not sourced and any initial condition in the gravitational potential is washed out as a^{-1} during the inflationary stage.

Comment: We might have computed the spectral index of the spectrum $\mathcal{P}_{\mathcal{R}}(k)$ by first solving the equation for the perturbation $\delta\phi_{\mathbf{k}}$ in a de Sitter stage, with $H = \text{constant}$ ($\epsilon = \eta = 0$), whose solution is Eq. (75) and then taking into account the time-evolution of the

Hubble rate and of ϕ introducing the subscript in H_k and $\dot{\phi}_k$. The time variation of the latter is determined by

$$\frac{d \ln \dot{\phi}_k}{d \ln k} = \left(\frac{d \ln \dot{\phi}_k}{dt} \right) \left(\frac{dt}{d \ln a} \right) \left(\frac{d \ln a}{d \ln k} \right) = \frac{\ddot{\phi}_k}{\dot{\phi}_k} \times \frac{1}{H} \times 1 = -\delta = \epsilon - \eta. \quad (171)$$

Correspondingly, $\dot{\phi}_k$ is the value of the time derivative of the inflaton field when a given wavelength $\sim k^{-1}$ crosses the horizon (from that point on the fluctuations remains frozen in). The curvature perturbation in such an approach would read

$$\mathcal{R}_{\mathbf{k}} \simeq \frac{H_k}{\dot{\phi}_k} \delta \phi_{\mathbf{k}} \simeq \frac{1}{2\pi} \left(\frac{H_k^2}{\dot{\phi}_k} \right).$$

Correspondingly

$$n_{\mathcal{R}} - 1 = \frac{d \ln \mathcal{P}_{\mathcal{R}}}{d \ln k} = \frac{d \ln H_k^4}{d \ln k} - \frac{d \ln \dot{\phi}_k^2}{d \ln k} = -4\epsilon + (2\eta - 2\epsilon) = 2\eta - 6\epsilon$$

which reproduces our previous findings.

During inflation the curvature perturbation is generated on superhorizon scales with a spectrum which is nearly scale invariant, that is is nearly independent from the wavelength $\lambda = \pi/k$: the amplitude of the fluctuation on superhorizon scales does not (almost) depend upon the time at which the fluctuations crosses the horizon and becomes frozen in. The small tilt of the power spectrum arises from the fact that the inflaton field is massive, giving rise to a nonvanishing η and because during inflation the Hubble rate is not exactly constant, but nearly constant, where ‘nearly’ is quantified by the slow-roll parameters ϵ .

Comment: From what found so far, we may conclude that on superhorizon scales the comoving curvature perturbation $\mathcal{R}$ and the uniform-density gauge curvature ζ satisfy on superhorizon scales the relation

$$\dot{\mathcal{R}}_{\mathbf{k}} \simeq \dot{\zeta}_{\mathbf{k}} \simeq 0.$$

An independent argument of the fact that they are nearly constant on superhorizon scales is given in the Appendix A.

7.12 Gauge-invariant computation of the curvature perturbation

In this subsection we would like to show how the computation of the curvature perturbation can be performed in a gauge-invariant way. We first rewrite Einstein’s equations in terms of Bardeen’s potentials (160) and (161)

$$\delta G_0^0 = \frac{2}{a^2} \left(-3 \mathcal{H} (\mathcal{H} \Phi + \Psi') + \nabla^2 \Psi + 3 \mathcal{H} (-\mathcal{H}' + \mathcal{H}^2) \left(\frac{E'}{2} - B \right) \right), \quad (172)$$

$$\delta G_i^0 = \frac{2}{a^2} \partial_i \left(\mathcal{H} \Phi + \Psi' + (\mathcal{H}' - \mathcal{H}^2) \left(\frac{E'}{2} - B \right) \right), \quad (173)$$

$$\begin{aligned}\delta G_j^i &= -\frac{2}{a^2} \left(\left((2\mathcal{H}' + 2\mathcal{H}^2) \Phi + \mathcal{H} \Phi' + \Psi'' + 2\mathcal{H} \Psi' + \frac{1}{2} \nabla^2 D \right) \delta_j^i \right. \\ &\quad \left. + \left(\mathcal{H}'' - \mathcal{H} \mathcal{H}' - \mathcal{H}^3 \right) \left(\frac{E'}{2} - B \right) \delta_j^i - \frac{1}{2} \partial^i \partial_j D \right),\end{aligned}\quad (174)$$

with $D = \Phi - \Psi$. These quantities are not gauge-invariant, but using the gauge transformations described in subsection 7.6, we can easily generalize them to gauge-invariant quantities

$$\delta G_0^{(\text{GI})0} = \delta G_0^0 + (G_0^0)' \left(\frac{E'}{2} - B \right), \quad (175)$$

$$\delta G_i^{(\text{GI})0} = \delta G_i^0 + \left(G_i^0 - \frac{1}{3} T_k^k \right) \partial_i \left(\frac{E'}{2} - B \right), \quad (176)$$

$$\delta G_j^{(\text{GI})i} = \delta G_j^i + (G_j^i)' \left(\frac{E'}{2} - B \right) \quad (177)$$

and

$$\delta T_0^{(\text{GI})0} = \delta T_0^0 + (T_0^0)' \left(\frac{E'}{2} - B \right) = -\delta \rho^{(\text{GI})}, \quad (178)$$

$$\delta T_i^{(\text{GI})0} = \delta T_i^0 + \left(T_i^0 - \frac{1}{3} T_k^k \right) \partial_i \left(\frac{E'}{2} - B \right) = (\rho + p) a^{-1} \delta v_i^{(\text{GI})}, \quad (179)$$

$$\delta T_j^{(\text{GI})i} = \delta T_j^i + (T_j^i)' \left(\frac{E'}{2} - B \right) = \delta p^{(\text{GI})} \quad (180)$$

where we have written the stress energy-momentum tensor as $T^{\mu\nu} = (\rho + p) u^\mu u^\nu + p \eta^{\mu\nu}$ with $u^\mu = (1, v^i)$.

Einstein's equations can now be written in a gauge-invariant way

$$- 3\mathcal{H}(\mathcal{H}\Phi + \Psi') + \nabla^2 \Psi \quad (181)$$

$$\begin{aligned}&= 4\pi G \left(-\Phi \phi'^2 + \delta\phi^{(\text{GI})} \phi' + \delta\phi^{(\text{GI})} \frac{\partial V}{\partial \phi} a^2 \right), \\ &\partial_i (\mathcal{H}\Phi + \Psi') = 4\pi G \left(\partial_i \delta\phi^{(\text{GI})} \phi' \right), \\ &\left((2\mathcal{H}' + \mathcal{H}^2) \Phi + \mathcal{H} \Phi' + \Psi'' + 2\mathcal{H} \Psi' + \frac{1}{2} \nabla^2 D \right) \delta_j^i - \frac{1}{2} \partial^i \partial_j D, \\ &= -4\pi G \left(\Phi \phi'^2 - \delta\phi^{(\text{GI})} \phi' + \delta\phi^{(\text{GI})} \frac{\partial V}{\partial \phi} a^2 \right) \delta_j^i.\end{aligned}\quad (182)$$

Taking $i \neq j$ from the third equation, we find $D = 0$, that is $\Psi = \Phi$ and from now on we can work with only the variable Φ . Using the background relation

$$2\left(\frac{a'}{a}\right)^2 - \frac{a''}{a} = 4\pi G \phi'^2 \quad (183)$$

we can rewrite the system of Eqs. (182) in the form

$$\begin{aligned}
\nabla^2 \Phi - 3 \mathcal{H} \Phi' - (\mathcal{H}' + 2 \mathcal{H}^2) &= 4 \pi G \left(\delta\phi^{(\text{GI})} \phi' + \delta\phi^{(\text{GI})} \frac{\partial V}{\partial \phi} a^2 \right) ; \\
\Phi' + \mathcal{H} \Phi &= 4 \pi G \left(\delta\phi^{(\text{GI})} \phi' \right) ; \\
\Phi'' + 3 \mathcal{H} \Phi' + (\mathcal{H}' + 2 \mathcal{H}^2) \Phi &= 4 \pi G \left(\delta\phi^{(\text{GI})} \phi' - \delta\phi^{(\text{GI})} \frac{\partial V}{\partial \phi} a^2 \right) .
\end{aligned} \tag{184}$$

Subtracting the first equation from the third, using the second equation to express $\delta\phi^{(\text{GI})}$ as a function of Φ and Φ' and using the Klein-Gordon equation one finally finds the

$$\Phi'' + 2 \left(\mathcal{H} - \frac{\phi''}{\phi'} \right) \Phi' - \nabla^2 \Phi + 2 \left(\mathcal{H}' - \mathcal{H} \frac{\phi''}{\phi'} \right) \Phi = 0 , \tag{185}$$

for the gauge-invariant potential Φ .

We now introduce the gauge-invariant quantity

$$u \equiv a \delta\phi^{(\text{GI})} + z \Psi , \tag{186}$$

$$z \equiv a \frac{\phi'}{\mathcal{H}} = a \frac{\dot{\phi}}{H} . \tag{187}$$

Notice that the variable u is equal to $-aQ$, the gauge-invariant inflaton perturbation on spatially flat gauges.

Eq. (185) becomes

$$u'' - \nabla^2 u - \frac{z''}{z} u = 0 , \tag{188}$$

while the two remaining equations of the system (184) can be written as

$$\nabla^2 \Phi = 4 \pi G \frac{\mathcal{H}}{a^2} (z u' - z' u) , \tag{189}$$

$$\left(\frac{a^2 \Phi}{\mathcal{H}} \right)' = 4 \pi G z u , \tag{190}$$

which allow to determine the variables Φ and $\delta\phi^{(\text{GI})}$.

We have now to solve Eq. (188). First, we have to evaluate $\frac{z''}{z}$ in terms of the slow-roll parameters

$$\frac{z'}{\mathcal{H}z} = \frac{a'}{\mathcal{H}a} + \frac{\phi''}{\mathcal{H}\phi'} - \frac{\mathcal{H}'}{\mathcal{H}^2} = \epsilon + \frac{\phi''}{\mathcal{H}\phi'} .$$

We then deduce that

$$\delta \equiv 1 - \frac{\phi''}{\mathcal{H}\phi'} = 1 + \epsilon - \frac{z'}{\mathcal{H}z} .$$

Keeping the slow-roll parameters constant in time (as we have mentioned, this corresponds to expand all quantities to first-order in the slow-roll parameters), we find

$$0 \simeq \delta' = \epsilon' (\simeq 0) - \frac{z''}{\mathcal{H}z} + \frac{z' \mathcal{H}'}{z \mathcal{H}^2} + \frac{(z')^2}{\mathcal{H}z^2},$$

from which we deduce

$$\frac{z''}{z} \simeq \frac{z' \mathcal{H}'}{z \mathcal{H}} + \frac{(z')^2}{z^2}.$$

Expanding in slow-roll parameters we find

$$\frac{z''}{z} \simeq (1 + \epsilon - \delta) (1 - \epsilon) \mathcal{H}^2 + (1 + \epsilon - \delta)^2 \mathcal{H}^2 \simeq \mathcal{H}^2 (2 + 2\epsilon - 3\delta).$$

If we set

$$\frac{z''}{z} = \frac{1}{\tau^2} \left(\nu^2 - \frac{1}{4} \right),$$

this corresponds to

$$\nu \simeq \frac{1}{2} \left[1 + 4 \frac{(1 + \epsilon - \delta)(2 - \delta)}{(1 - \epsilon)^2} \right]^{1/2} \simeq \frac{3}{2} + (2\epsilon - \delta) \simeq \frac{3}{2} + 3\epsilon - \eta.$$

On subhorizon scales ($k \gg aH$), the solution of equation (188) is obviously $u_{\mathbf{k}} \simeq e^{-ik\tau} / \sqrt{2k}$. Rewriting Eq. (190) as

$$\Phi_{\mathbf{k}} = -\frac{4\pi G a^2}{k^2} \frac{\dot{\phi}^2}{H} \left(\frac{H}{a\dot{\phi}} u_{\mathbf{k}} \right),$$

we infer that on subhorizon scales

$$\Phi_{\mathbf{k}} \simeq i \frac{4\pi G \dot{\phi}}{\sqrt{2k^3}} e^{-i\frac{k}{a}}.$$

On superhorizon scales ($k \ll aH$), one obvious solution to Eq. (188) is $u_{\mathbf{k}} \propto z$. To find the other solution, we may set $u_{\mathbf{k}} = z \tilde{u}_{\mathbf{k}}$, which satisfies the equation

$$\frac{\tilde{u}_{\mathbf{k}}''}{\tilde{u}_{\mathbf{k}}'} = -2 \frac{z'}{z},$$

which gives

$$\tilde{u}_{\mathbf{k}} = \int^{\tau} \frac{d\tau'}{z^2(\tau')}.$$

On superhorizon scales therefore we find

$$u_{\mathbf{k}} = c_1(k) \frac{a\dot{\phi}}{H} + c_2(k) \frac{a\dot{\phi}}{H} \int^t dt' \frac{H^2}{a^3 \dot{\phi}^2} \simeq c_1(k) \frac{a\dot{\phi}}{H} - c_2(k) \frac{1}{3a^2 \dot{\phi}},$$

where the last passage has been performed supposing a de Sitter epoch, $H = \text{constant}$. The first piece is the constant mode $c_1(k)z$, while the second is the decreasing mode. To find the constant $c_1(k)$, we apply what we have learned in subsection 6.5. We know that on superhorizon scales the exact solution of equation (188) is

$$u_{\mathbf{k}} = \frac{\sqrt{\pi}}{2} e^{i(\nu+\frac{1}{2})\frac{\pi}{2}} \sqrt{-\tau} H_\nu^{(1)}(-k\tau). \quad (191)$$

On superhorizon scales, since $H_\nu^{(1)}(x \ll 1) \sim \sqrt{2/\pi} e^{-i\frac{\pi}{2}} 2^{\nu-\frac{3}{2}} (\Gamma(\nu_\chi)/\Gamma(3/2)) x^{-\nu}$, the fluctuation (191) becomes

$$u_{\mathbf{k}} = e^{i(\nu-\frac{1}{2})\frac{\pi}{2}} 2^{(\nu-\frac{3}{2})} \frac{\Gamma(\nu)}{\Gamma(3/2)} \frac{1}{\sqrt{2k}} (-k\tau)^{\frac{1}{2}-\nu}.$$

Therefore

$$c_1(k) = \lim_{k \rightarrow 0} \left| \frac{u_{\mathbf{k}}}{z} \right| = \frac{H}{a\dot{\phi}} \frac{1}{\sqrt{2k}} \left(\frac{k}{aH} \right)^{\frac{1}{2}-\nu} = \frac{H}{\dot{\phi}} \frac{1}{\sqrt{2k^3}} \left(\frac{k}{aH} \right)^{\eta-3\epsilon} \quad (192)$$

The last steps consist in relating the variable u to the comoving curvature $\mathcal{R}$ and to the gravitational potential Φ . The comoving curvature takes the form

$$\mathcal{R} \equiv -\Psi - \frac{H}{\dot{\phi}'} \delta\phi^{(\text{GI})} = -\frac{u}{z}. \quad (193)$$

Since $z = a\dot{\phi}/H = a\sqrt{2\epsilon}m_{\text{Pl}}$, the power spectrum of the comoving curvature can be expressed on superhorizon scales as

$$\mathcal{P}_{\mathcal{R}}(k) = \frac{k^3}{2\pi^2} \left| \frac{u_{\mathbf{k}}}{z} \right|^2 = \frac{1}{2m_{\text{Pl}}^2\epsilon} \left(\frac{H}{2\pi} \right)^2 \left(\frac{k}{aH} \right)^{n_{\mathcal{R}}-1} \equiv A_{\mathcal{R}}^2 \left(\frac{k}{aH} \right)^{n_{\mathcal{R}}-1} \quad (194)$$

with

$$n_{\mathcal{R}} - 1 = 3 - 2\nu = 2\eta - 6\epsilon. \quad (195)$$

These results reproduce those found in the previous subsection.

The last step is to find the behaviour of the gauge-invariant potential Φ on superhorizon scales. If we recast equation (190) in the form

$$u_{\mathbf{k}} = \frac{1}{4\pi G} \frac{H}{\dot{\phi}} \left(\frac{a}{H} \Phi_{\mathbf{k}} \right), \quad (196)$$

we can infer that on superhorizon scales the nearly constant mode of the gravitational potential during inflation reads

$$\Phi_{\mathbf{k}} = c_1(k) \left[1 - \frac{H}{a} \int^t dt' a(t') \right] \simeq -c_1(k) \frac{\dot{H}}{H^2} = \epsilon c_1(k) \simeq \epsilon \frac{u_{\mathbf{k}}}{z} \simeq -\epsilon \mathcal{R}_{\mathbf{k}}. \quad (197)$$

Indeed, plugging this solution into Eq. (196), one reproduces $u_{\mathbf{k}} = c_1(k) \frac{a\dot{\phi}}{H}$.

7.13 Gravitational waves

Quantum fluctuations in the gravitational fields are generated in a similar fashion of that of the scalar perturbations discussed so far. A gravitational wave may be viewed as a ripple of spacetime in the FRW background metric (1) and in general the linear tensor perturbations may be written as

$$g_{\mu\nu} = a^2(\tau) \left[-d\tau^2 + (\delta_{ij} + h_{ij}) dx^i dx^j \right],$$

where $|h_{ij}| \ll 1$. The tensor h_{ij} has six degrees of freedom, but, as we studied in subsection 7.1, the tensor perturbations are traceless, $\delta^{ij} h_{ij} = 0$, and transverse $\partial^i h_{ij} = 0$ ($i = 1, 2, 3$). With these 4 constraints, there remain 2 physical degrees of freedom, or polarizations, which are usually indicated $\lambda = +, \times$. More precisely, we can write

$$h_{ij} = h_+ e_{ij}^+ + h_\times e_{ij}^\times,$$

where e^+ and $e^\times$ are the polarization tensors which have the following properties

$$\begin{aligned} e_{ij} &= e_{ji}, \quad k^i e_{ij} = 0, \quad e_{ii} = 0, \\ e_{ij}(-\mathbf{k}, \lambda) &= e_{ij}^*(\mathbf{k}, \lambda), \quad \sum_\lambda e_{ij}^*(\mathbf{k}, \lambda) e^{ij}(\mathbf{k}, \lambda) = 4. \end{aligned}$$

Notice also that the tensors h_{ij} are gauge-invariant and therefore represent physical degrees of freedom.

If the stress-energy momentum tensor is diagonal, as the one provided by the inflaton potential $T_{\mu\nu} = \partial_\mu \phi \partial_\nu \phi - g_{\mu\nu} \mathcal{L}$, the tensor modes do not have any source in their equation and their action can be written as

$$\frac{m_{\text{Pl}}^2}{2} \int d^4x \sqrt{-g} \frac{1}{2} \partial_\sigma h_{ij} \partial^\sigma h_{ij},$$

that is the action of four independent massless scalar fields. The gauge-invariant tensor amplitude

$$v_{\mathbf{k}} = a m_{\text{Pl}} \frac{1}{\sqrt{2}} h_{\mathbf{k}},$$

satisfies therefore the equation

$$v_{\mathbf{k}}'' + \left(k^2 - \frac{a''}{a} \right) v_{\mathbf{k}} = 0,$$

which is the equation of motion of a massless scalar field in a quasi-de Sitter epoch. We can therefore make use of the results present in subsection 6.5 and Eq. (87) to conclude that on superhorizon scales the tensor modes scale like

$$|v_{\mathbf{k}}| = \left(\frac{H}{2\pi} \right) \left(\frac{k}{aH} \right)^{\frac{3}{2} - \nu_T},$$

where

$$\nu_T \simeq \frac{3}{2} - \epsilon.$$

Since fluctuations are (nearly) frozen in on superhorizon scales, a way of characterizing the tensor perturbations is to compute the spectrum on scales larger than the horizon

$$\mathcal{P}_T(k) = \frac{k^3}{2\pi^2} \sum_{\lambda} |h_{\mathbf{k}}|^2 = 4 \times 2 \frac{k^3}{2\pi^2} |v_{\mathbf{k}}|^2. \quad (198)$$

This gives the power spectrum on superhorizon scales

$$\mathcal{P}_T(k) = \frac{8}{m_{\text{Pl}}^2} \left(\frac{H}{2\pi} \right)^2 \left(\frac{k}{aH} \right)^{n_T} \equiv A_T^2 \left(\frac{k}{aH} \right)^{n_T}$$

where we have defined the *spectral index* n_T of the tensor perturbations as

$$n_T = \frac{d \ln \mathcal{P}_T}{d \ln k} = 3 - 2\nu_T = -2\epsilon.$$

The tensor perturbation is almost scale-invariant. Notice that the amplitude of the tensor modes depends only on the value of the Hubble rate during inflation. This amounts to saying that it depends only on the energy scale $V^{1/4}$ associated to the inflaton potential. A detection of gravitational waves from inflation will be therefore a direct measurement of the energy scale associated to inflation.

7.14 The consistency relation

The results obtained so far for the scalar and tensor perturbations allow to predict a *consistency relation* which holds for the models of inflation addressed in these lectures, *i.e.* the models of inflation driven by one-single field ϕ . We define tensor-to-scalar amplitude ratio to be

$$r = \frac{\frac{1}{100} A_T^2}{\frac{4}{25} A_{\mathcal{R}}^2} = \frac{\frac{1}{100} 8 \left(\frac{H}{2\pi m_{\text{Pl}}} \right)^2}{\frac{4}{25} (2\epsilon)^{-1} \left(\frac{H}{2\pi m_{\text{Pl}}} \right)^2} = \epsilon.$$

This means that

$$r = -\frac{n_T}{2}$$

One-single models of inflation predict that during inflation driven by a single scalar field, the ratio between the amplitude of the tensor modes and that of the curvature perturbations is equal to minus one-half of the tilt of the spectrum of tensor modes. If this relation turns out to be falsified by the future measurements of the CMB anisotropies, this does not mean that inflation is wrong, but only that inflation has not been driven by only one field. Generalizations to two-field models of inflation can be found for instance in Refs. [10, 12].

8 The post-inflationary evolution of the cosmological perturbations

So far, we have computed the evolution of the cosmological perturbations within the horizon and outside the horizon during inflation. However, what we are really interested in is their evolution after inflation and to compute the amplitude of perturbations when they re-enter the horizon during radiation- or matter-domination.

To this purpose, we use the following procedure. We use Eqs. (177) and (180) to write

$$\nabla^2 \Phi - 3 \mathcal{H} \Phi' - (\mathcal{H}' + 2 \mathcal{H}^2) = 4 \pi G a^2 \delta \rho^{(\text{GI})}; \quad (199)$$

$$\Phi' + \mathcal{H} \Phi = 4 \pi G a^2 (\rho + p) \delta v^{(\text{GI})}; \quad (200)$$

$$\Phi'' + 3 \mathcal{H} \Phi' + (\mathcal{H}' + 2 \mathcal{H}^2) \Phi = 4 \pi G a^2 \delta p^{(\text{GI})}. \quad (201)$$

Combining these equations one finds

$$\Phi'' + 3 \mathcal{H} (1 + c_s^2) \Phi' - c_s^2 \nabla^2 \Phi + [2 \mathcal{H}' + (1 + c_s^2) \mathcal{H}^2] \Phi = 0, \quad (202)$$

where $c_s^2 = \dot{p}/\dot{\rho}$. This equation can be rewritten as

$$\dot{\mathcal{R}}_{\mathbf{k}} = 0$$

where we have set

$$\mathcal{R}_{\mathbf{k}} = -\Phi - \frac{2}{3} \frac{\mathcal{H}^{-1} \dot{\Phi} + \Phi}{(1 + w)} - \Phi.$$

Here $w = p/\rho$. Notice that during inflation, when $p = \frac{1}{2} \dot{\phi}^2 - V$ and $\rho = \frac{1}{2} \dot{\phi}^2 + V$, $\mathcal{R}_{\mathbf{k}}$ takes the form

$$\mathcal{R}_{\mathbf{k}} = -\Phi_{\mathbf{k}} - \frac{1}{\epsilon \mathcal{H}} (\phi'_{\mathbf{k}} + \mathcal{H} \Phi_{\mathbf{k}}), \quad (203)$$

which using the equation

$$\Phi' + \mathcal{H} \Phi = 4 \pi G (\delta \Phi^{(\text{GI})} \phi')$$

reduces to the comoving curvature perturbation (193).

Eq. (202) is solved by

$$\Phi_{\mathbf{k}} = c_1(k) \left(1 - \frac{\mathcal{H}}{a^2} \int^\tau d\tau' a^2(\tau') \right) + c_2(k) \frac{\mathcal{H}}{a^2} \quad (\text{ON SUPERHORIZON SCALES})$$

This nearly constant solution can be rewritten in cosmic time as

$$\Phi_{\mathbf{k}} = c_1(k) \left(1 - \frac{H}{a} \int^t dt' a(t') \right).$$

which is the same form of solution we found during inflation, see Eq. (197). This explains why we can choose the constant $c_1(k)$ to be the one given by Eq. (192), $c_1(k) = |u_{\mathbf{k}}/z|$ for superhorizon scales.

Since during radiation-domination $a \sim t^n$ with $n = 1/2$ and during matter-domination $a \sim t^n$ with $n = 2/3$, we can write

$$\Phi_{\mathbf{k}} = c_1(k) \left(1 - \frac{H}{a} \int^t dt' a(t') \right) = c_1(k) \left(1 - \frac{n}{n+1} \right) = \frac{c_1(k)}{n+1} = \mathcal{R}_{\mathbf{k}} \begin{cases} \frac{2}{3} & \text{RD} \\ \frac{3}{5} & \text{MD} \end{cases}$$

This relation tells us that, after a gravitational perturbation with a given wavelength is generated during inflation, it evolves on superhorizon scales after inflation simply slightly rescaling its amplitude. When the given wavelength re-enters the horizon, the amplitude of the gravitational potential depends upon the time of re-enter. If the perturbation re-enters the horizon when the universe is still dominated by radiation, then $\Phi_{\mathbf{k}} = \frac{2}{3}\mathcal{R}_{\mathbf{k}}$; if the perturbation re-enters the horizon when the universe is dominated by matter, then $\Phi_{\mathbf{k}} = \frac{3}{5}\mathcal{R}_{\mathbf{k}}$. For instance, the power spectrum of the gravitational perturbations during matter-domination reads

$$\mathcal{P}_{\Phi} = \left(\frac{3}{5} \right)^2 \mathcal{P}_{\mathcal{R}} = \left(\frac{3}{5} \right)^2 \frac{1}{2m_{\text{Pl}}^2 \epsilon} \left(\frac{H}{2\pi} \right)^2 \left(\frac{k}{aH} \right)^{n_{\mathcal{R}}-1}$$

8.1 From the inflationary seeds to the matter power spectrum

As the curvature perturbations enter the causal horizon during radiation- or matter-domination, they create density fluctuations $\delta\rho_{\mathbf{k}}$ via gravitational attractions of the potential wells. The density contrast $\delta_{\mathbf{k}} = \frac{\delta\rho_{\mathbf{k}}}{\bar{\rho}}$ can be deduced from Poisson equation

$$\frac{k^2}{a^2} = -4\pi G \delta\rho_{\mathbf{k}} = -4\pi G \frac{\delta\rho_{\mathbf{k}}}{\bar{\rho}} \bar{\rho} = \frac{3}{2} H^2 \frac{\delta\rho_{\mathbf{k}}}{\bar{\rho}}$$

where $\bar{\rho}$ is the background average energy density. This means that

$$\delta_{\mathbf{k}} = \frac{2}{3} \left(\frac{k}{aH} \right)^2 \Phi_{\mathbf{k}}.$$

From this expression we can compute the power spectrum of matter density perturbations induced by inflation when they re-enter the horizon during matter-domination

$$\mathcal{P}_{\delta\rho} = \langle |\delta_{\mathbf{k}}|^2 \rangle = A \left(\frac{k}{aH} \right)^n = \frac{2\pi^2}{k^3} \left(\frac{2}{5} \right)^2 A_{\mathcal{R}}^2 \left(\frac{k}{aH} \right)^4 \left(\frac{k}{aH} \right)^{n_{\mathcal{R}}-1}$$

from which we deduce that matter perturbations scale linearly with the wavenumber and have a scalar tilt

$$n = n_{\mathcal{R}} = 1 + 2\eta - 6\epsilon.$$

The primordial spectrum $\mathcal{P}_{\delta\rho}$ is of course reprocessed by gravitational instabilities after the universe becomes matter-dominated. Indeed, as we have seen in section 6, perturbations evolve after entering the horizon and the power spectrum will not remain constant. To see how the density contrast is reprocessed we have first to analyze how it evolves on superhorizon scales before horizon-crossing. We use the following trick. Consider a flat universe with average energy density $\bar{\rho}$. The corresponding Hubble rate is

$$H^2 = \frac{8\pi G}{3} \bar{\rho}.$$

A small positive fluctuation $\delta\rho$ will cause the universe to be closed

$$H^2 = \frac{8\pi G}{3} (\bar{\rho} + \delta\rho) - \frac{k}{a^2}.$$

Subtracting the two equations we find

$$\frac{\delta\rho}{\rho} = \frac{3}{8\pi G} \frac{k}{a^2 \rho} \sim \begin{cases} a^2 & \text{RD} \\ a & \text{MD} \end{cases}$$

Notice that $\Phi_{\mathbf{k}} \sim \delta\rho a^2/k^2 \sim (\delta\rho/\rho)\rho a^2/k^2 = \text{constant}$ for both RD and MD which confirms our previous findings.

When the matter densities enter the horizon, they do not increase appreciably before matter-domination because before matter-domination pressure is too large and does not allow the matter inhomogeneities to grow. On the other hand, the suppression of growth due to radiation is restricted to scales smaller than the horizon, while large-scale perturbations remain unaffected. This is the reason why the horizon size at equality sets an important scale for structure growth

$$k_{\text{EQ}} = H^{-1}(a_{\text{EQ}}) \simeq 0.08 h \text{ Mpc}^{-1}.$$

Therefore, perturbations with $k \gg k_{\text{EQ}}$ are perturbations which have entered the horizon before matter-domination and have remained nearly constant till equality. This means that they are suppressed with respect to those perturbations having $k \ll k_{\text{EQ}}$ by a factor $(a_{\text{ENT}}/a_{\text{EQ}})^2 = (k_{\text{EQ}}/k)^2$. If we define the transfer function $T(k)$ by the relation $\mathcal{R}_{\text{final}} = T(k) \mathcal{R}_{\text{initial}}$ we find therefore that roughly speaking

$$T(k) = \begin{cases} 1 & k \ll k_{\text{EQ}}, \\ (k_{\text{EQ}}/k)^2 & k \gg k_{\text{EQ}}. \end{cases}$$

The corresponding power spectrum will be

$$\mathcal{P}_{\delta\rho}(k) \sim \begin{cases} \left(\frac{k}{aH}\right) & k \ll k_{\text{EQ}}, \\ \left(\frac{k}{aH}\right)^{-3} & k \gg k_{\text{EQ}}. \end{cases}$$

Of course, a more careful computation needs to include many other effects such as neutrino free-streaming, photon diffusion and the diffusion of baryons along with photons. It is encouraging however that this rough estimate turns out to be confirmed by present data on large scale structures [30].

8.2 From inflation to large-angle CMB anisotropy

Temperature fluctuations in the CMB arise due to five distinct physical effects: our peculiar velocity with respect to the cosmic rest frame; fluctuations in the gravitational potential on the last scattering surface; fluctuations intrinsic to the radiation field itself on the last-scattering surface; the peculiar velocity of the last-scattering surface and damping of anisotropies if the universe should be re-ionized after decoupling. The first effect gives rise to the dipole anisotropy. The second effect, known as the Sachs-Wolfe effect is the dominant contribution to the anisotropy on large-angular scales, $\theta \gg \theta_{\text{HOR}} \sim 1^\circ$. The last three effects provide the dominant contributions to the anisotropy on small angular scales, $\theta \ll 1^\circ$.

We consider here the temperature fluctuations on large-angular scales that arise due to the Sachs-Wolfe effect. These anisotropies probe length scales that were superhorizon sized at photon decoupling and therefore insensitive to microphysical processes. On the contrary, they provide a probe of the original spectrum of primeval fluctuations produced during inflation.

To proceed, we consider the CMB anisotropy measured at positions other than our own and at earlier times. This is called the brightness function $\Theta(t, \mathbf{x}, \mathbf{n}) \equiv \delta T(t, \mathbf{x}, \mathbf{n})/T(t)$. The photons with momentum $\mathbf{p}$ in a given range d^3p have intensity I proportional to $T^4(t, \mathbf{x}, \mathbf{n})$ and therefore $\delta I/I = 4\Theta$. The brightness function depends upon the direction $\mathbf{n}$ of the photon momentum or, equivalently, on the direction of observation $\mathbf{e} = -\mathbf{n}$. Because the CMB travels freely from the last-scattering, we can write

$$\frac{\delta T}{T} = \Theta(t_{\text{LS}}, \mathbf{x}_{\text{LS}}, \mathbf{n}) + \left(\frac{\delta T}{T}\right)_*,$$

where $\mathbf{x}_{\text{LS}} = -x_{\text{LS}}\mathbf{n}$ is the point of the origin of the photon coming from the direction $\mathbf{e}$. The comoving distance of the last-scattering distance is $x_{\text{LS}} = 2/H_0$. The first term corresponds to the anisotropy already present at last scattering and the second term is the additional anisotropy acquired during the travel towards us, equal to minus the fractional perturbation in the redshift of the radiation. Notice that the separation between each term depends on the slicing, but the sum is not.

Consider the redshift perturbation on comoving slicing. We imagine the universe populated by comoving observers along the line of sight. The relative velocity of adjacent comoving observers is equal to their distance times the velocity gradient measured along $\mathbf{n}$ of the photon. In the unperturbed universe, we have $\mathbf{u} = H\mathbf{r}$, leading to the velocity gradient $u_{ij} = \partial u_i / \partial r_j = u_{ij} = H(t)\delta_{ij}$ with zero vorticity and shear. Including a peculiar velocity

field as perturbation, $\mathbf{u} = H\mathbf{r} + \mathbf{v}$ and $u_{ij} = H(t)\delta_{ij} + \frac{1}{a}\frac{\partial v_i}{\partial x_j}$. The corresponding Doppler shift is

$$\frac{d\lambda}{\lambda} = \frac{da}{a} + n_i n_j \frac{\partial v_i}{\partial x_j} dx.$$

The perturbed FRW equation is

$$\delta H = \frac{1}{3}\nabla \cdot \mathbf{v},$$

while

$$(\delta\rho)^\cdot = -3\rho\delta H - 3H\delta\rho.$$

Instead of $\delta\rho$, let us work with the density contrast $\delta = \delta\rho/\rho$. Remembering that $\rho \sim a^{-3}$, we find that $\dot{\delta} = -3\delta H$, which give

$$\nabla \cdot \mathbf{v} = -\dot{\delta}.$$

From Euler equation $\dot{\mathbf{u}} = -\rho^{-1}\nabla p - \nabla\Phi$, we deduce $\dot{\mathbf{v}} + H\mathbf{v} = -\nabla\Phi - \rho^{-1}\nabla p$. Therefore, for $a \sim t^{2/3}$ and negligible pressure gradient, since the gravitational potential is constant, we find

$$\mathbf{v} = -t\nabla\Phi$$

leading to

$$\left(\frac{\delta T}{T}\right)_* = \int_0^{x_{\text{LS}}} \frac{t}{a} \frac{d^2\Phi}{dx^2} dx. \quad (204)$$

The photon trajectory is $a d\mathbf{x}/dt = \mathbf{n}$. Using $a \sim t^{2/3}$ gives

$$x(t) = \int_t^{t_0} \frac{dt'}{a} = 3 \left(\frac{a_0}{t_0} - \frac{t}{a} \right).$$

Integrating by parts Eq. (204), we finally find

$$\left(\frac{\delta T}{T}\right)_* = \frac{1}{3} [\Phi(\mathbf{x}_{\text{LS}}) - \Phi(0)] + \mathbf{e} \cdot [\mathbf{v}(0, t_0) - \mathbf{v}(\mathbf{x}_{\text{LS}}, t_{\text{LS}})].$$

The potential at our position contributes only to the unobservable monopole and can be dropped. On scales outside the horizon, $\mathbf{v} = -t\nabla\Phi \sim 0$. The remaining term is the Sachs-Wolfe effect

$$\frac{\delta T(\mathbf{e})}{T} = \frac{1}{3}\Phi(\mathbf{x}_{\text{LS}}) = \frac{1}{5}\mathcal{R}(\mathbf{x}_{\text{LS}}).$$

Therefore, at large angular scales, the theory of cosmological perturbations predicts a remarkable simple formula relating the CMB anisotropy to the curvature perturbation generated during inflation.

In section 3, we have seen that the temperature anisotropy is commonly expanded in spherical harmonics $\frac{\Delta T}{T}(x_0, \tau_0, \mathbf{n}) = \sum_{\ell m} a_{\ell m}(x_0) Y_{\ell m}(\mathbf{n})$, where x_0 and τ_0 are our position and the preset time, respectively, $\mathbf{n}$ is the direction of observation, ℓ 's are the different multipoles and $\langle a_{\ell m} a_{\ell' m'}^* \rangle = \delta_{\ell \ell'} \delta_{m m'} C_\ell$, where the deltas are due to the fact that the process that created the anisotropy is statistically isotropic. The C_ℓ are the so-called CMB power spectrum. For homogeneity and isotropy, the C_ℓ 's are neither a function of x_0 , nor of m . The two-point-correlation function is related to the C_ℓ 's according to Eq. (37).

For adiabatic perturbations we have seen that on large scales, larger than the horizon on the last-scattering surface (corresponding to angles larger than $\theta_{\text{HOR}} \sim 1^\circ$) $\delta T/T = \frac{1}{3} \Phi(\mathbf{x}_{\text{LS}})$. In Fourier transform

$$\frac{\delta T(\mathbf{k}, \tau_0, \mathbf{n})}{T} = \frac{1}{3} \Phi_{\mathbf{k}} e^{i \mathbf{k} \cdot \mathbf{n} (\tau_0 - \tau_{\text{LS}})} \quad (205)$$

Using the decomposition

$$\exp(i \mathbf{k} \cdot \mathbf{n} (\tau_0 - \tau_{\text{LS}})) = \sum_{\ell=0}^{\infty} (2\ell + 1) i^\ell j_\ell(k(\tau_0 - \tau_{\text{LS}})) P_\ell(\mathbf{k} \cdot \mathbf{n}) \quad (206)$$

where j_ℓ is the spherical Bessel function of order ℓ and substituting, we get

$$\begin{aligned} & \left\langle \frac{\delta T(x_0, \tau_0, \mathbf{n})}{T} \frac{\delta T(x_0, \tau_0, \mathbf{n}')}{T} \right\rangle = \\ &= \frac{1}{V} \int d^3 x \left\langle \frac{\delta T(x_0, \tau_0, \mathbf{n})}{T} \frac{\delta T(x_0, \tau_0, \mathbf{n}')}{T} \right\rangle = \\ &= \frac{1}{(2\pi)^3} \int d^3 k \left\langle \frac{\delta T(\mathbf{k}, \tau_0, \mathbf{n})}{T} \left(\frac{\delta T(\mathbf{k}, \tau_0, \mathbf{n}')}{T} \right)^* \right\rangle = \\ &= \frac{1}{(2\pi)^3} \int d^3 k \left(\left\langle \frac{1}{3} |\Phi|^2 \right\rangle \sum_{\ell, \ell'=0}^{\infty} (2\ell + 1)(2\ell' + 1) j_\ell(k(\tau_0 - \tau_{\text{LS}})) \right. \\ & \quad \left. j_{\ell'}(k(\tau_0 - \tau_{\text{LS}})) P_\ell(\mathbf{k} \cdot \mathbf{n}) P_{\ell'}(\mathbf{k}' \cdot \mathbf{n}') \right) \end{aligned} \quad (207)$$

Inserting $P_\ell(\mathbf{k} \cdot \mathbf{n}) = \frac{4\pi}{2\ell+1} \sum_m Y_{\ell m}^*(\mathbf{k}) Y_{\ell m}(\mathbf{n})$ and analogously for $P_\ell(\mathbf{k}' \cdot \mathbf{n}')$, integrating over the directions $d\Omega_k$ generates $\delta_{\ell \ell'} \delta_{m m'} \sum_m Y_{\ell m}^*(\mathbf{n}) Y_{\ell m}(\mathbf{n}')$. Using as well $\sum_m Y_{\ell m}^*(\mathbf{n}) Y_{\ell m}(\mathbf{n}') = \frac{2\ell+1}{4\pi} P_\ell(\mathbf{n} \cdot \mathbf{n}')$, we get

$$\begin{aligned} & \left\langle \frac{\delta T(x_0, \tau_0, \mathbf{n})}{T} \frac{\delta T(x_0, \tau_0, \mathbf{n}')}{T} \right\rangle \\ &= \sum_\ell \frac{2\ell+1}{4\pi} P_\ell(\mathbf{n} \cdot \mathbf{n}') \frac{2}{\pi} \int \frac{dk}{k} \left\langle \frac{1}{9} |\Phi|^2 \right\rangle k^3 j_\ell^2(k(\tau_0 - \tau_{\text{LS}})). \end{aligned} \quad (208)$$

Comparing this expression with that for the C_ℓ , we get the expression for the C_ℓ^{AD} , where the suffix “AD” stays for adiabatic

$$C_\ell^{\text{AD}} = \frac{2}{\pi} \int \frac{dk}{k} \left\langle \frac{1}{9} |\Phi|^2 \right\rangle k^3 j_\ell^2(k(\tau_0 - \tau_{\text{LS}})) \quad (209)$$

which is valid for $2 \leq \ell \ll (\tau_0 - \tau_{\text{LS}})/\tau_{\text{LS}} \sim 100$.

If we generically indicate by $\langle |\Phi_{\mathbf{k}}|^2 \rangle k^3 = A^2 (k\tau_0)^{n-1}$, we can perform the integration and get

$$\frac{\ell(\ell+1)C_\ell^{\text{AD}}}{2\pi} = \left[\frac{\sqrt{\pi}}{2} \ell(\ell+1) \frac{\Gamma(\frac{3-n}{2})\Gamma(\ell + \frac{n-1}{2})}{\Gamma(\frac{4-n}{2})\Gamma(\ell + \frac{5-n}{2})} \right] \frac{A^2}{9} \left(\frac{H_0}{2} \right)^{n-1} \quad (211)$$

For $n \simeq 1$ and $\ell \gg 1$, we can approximate this expression to

$$\frac{\ell(\ell+1)C_\ell^{\text{AD}}}{2\pi} = \frac{A^2}{9}. \quad (212)$$

This result shows that inflation predicts a very flat spectrum for low ℓ . This prediction has been confirmed by the COBE satellite [49]. Furthermore, since inflation predicts $\Phi_{\mathbf{k}} = \frac{3}{5}\mathcal{R}_{\mathbf{k}}$, we find that

$$\pi \ell(\ell+1)C_\ell^{\text{AD}} = \frac{A_{\mathcal{R}}^2}{25} = \frac{1}{25} \frac{1}{2 m_{\text{Pl}}^2 \epsilon} \left(\frac{H}{2\pi} \right)^2. \quad (213)$$

COBE data imply that $\frac{\ell(\ell+1)C_\ell^{\text{AD}}}{2\pi} \simeq 10^{-10}$ or

$$\left(\frac{V}{\epsilon} \right)^{1/4} \simeq 6.7 \times 10^{16} \text{ GeV}$$

Take for instance a model of chaotic inflation with quadratic potential $V(\phi) = \frac{1}{2}m_\phi^2\phi^2$. Using Eq. (59) one easily computes that when there are ΔN e-foldings to go, the value of the inflaton field is $\phi_{\Delta N}^2 = (\Delta N/2\pi G)$ and the corresponding value of ϵ is $1/(2\Delta N)$. Taking $\Delta N \simeq 60$ (corresponding to large-angle CMB anisotropies), one finds that COBE normalization imposes $m_\phi \simeq 10^{13}$ GeV.

9 Conclusions

Along these lectures, we have learned that a stage of inflation during the early epochs of the evolution of the universe solves many drawbacks of the standard Big-Bang cosmology, such as the flatness or entropy problem and the horizon problem. Luckily, despite inflation occurs after a tiny bit after the bang, it leaves behind some observable predictions:

- *The universe should be flat*, that is the total density of all components of matter should sum to the critical energy density and $\Omega_0 = 1$. The current data on the CMB anisotropies offer a spectacular confirmation of such a prediction. The universe appears indeed to be spatially flat.
- *Primordial perturbations are adiabatic*. Inflation provides the seeds for the cosmological perturbations. In one-single field models of inflation, the perturbations are *adiabatic* or curvature perturbations, *i.e.* they are fluctuations in the total energy density of the universe or, equivalently, scalar perturbations to the spacetime metric. Adiabaticity

implies that the spatial distribution of each species in the universe is the same, that is the ratio of number densities of any two of these species is everywhere the same. Adiabatic perturbations predict a contribution to the CMB anisotropy which is related to the curvature perturbation $\mathcal{R}$ on large angles, $\delta T/T = \frac{1}{5} \mathcal{R}$, and are in excellent agreement with the CMB data. Adiabatic perturbations can be contrasted to isocurvature perturbations which are fluctuations in the ratios between the various species in the universe. Isocurvature perturbations predict that on large angles $\delta T/T = -2\Phi$ and are presently ruled out, even though a certain amount of isocurvature perturbations, possibly correlated with the adiabatic fluctuations, cannot be excluded by present CMB data [4].

- *Primordial perturbations are almost scale-independent.* The primordial power spectrum predicted by inflation has a characteristic feature, it is almost scale-independent, that is the spectral index $n_{\mathcal{R}}$ is very close to unity. Possible deviations from exact scale-independence arise because during inflation the inflaton is not massless and the Hubble rate is not exactly constant. A recent analysis [14] shows that $n_{\mathcal{R}} = 0.97^{+0.08}_{-0.05}$, again in agreement with the theoretical prediction.
- *Primordial perturbations are nearly gaussian.* The fact that cosmological perturbations are tiny allow their analysis in terms of linear perturbation theory. Non-gaussian features are therefore suppressed since the non-linearities of the inflaton potential and of the metric perturbations are suppressed. Non-gaussian features are indeed present, but may appear only at the second-order in deviations from the homogeneous background solution and are therefore small [2]. This is rigously true only for one-single field models of inflation. Many-field models of inflation may give rise to some level of non-gaussianity [11]. If the next generation of satellites will detect a non-negligible amount of non-gaussianity in the CMB anisotropy, this will rule out one-single field models of inflation.
- *Production of gravitational waves.* A stochastic background of gravitational waves is produced during inflation in the very same way classical perturbations to the inflaton field are generated. The spectrum of such gravitational waves is again flat, *i.e.* scale-independent and the tensor-to-scalar amplitude ratio r is, at least in one-single models of inflation, related to the spectral index n_T by the consistency relation $r = -n_T/2$. A confirmation of such a relation would be a spectacular proof of one-single field models of inflation. The detection of the primordial stochastic background of gravitational waves from inflation is challenging, but would not only set the energy scale of inflation, but would also give the opportunity of discriminating among different models of inflation [33, 5].

Acknowledgments

The author would like to thank the organizers of the School, G. Dvali, A. Perez-Lorezana, G. Senjanovic, G. Thompson and F. Vissani for providing the students and the lecturers with such an excellent and stimulating environment. He also thanks the students for their questions and enthusiasm.

A Evolution of the curvature perturbation on super-horizon scales

In this appendix, we repeat the general arguments following from energy-momentum conservation given in Ref. [66] to show that the curvature perturbation on constant-time hypersurfaces ψ is constant on superhorizon scales if perturbations are adiabatic.

The constant-time hypersurfaces are orthogonal to the unit time-like vector field $n^\mu = (1 - A, -\partial^i B/2)$. Local conservation of the energy-momentum tensor tells us that $T^\mu_{\nu;\mu} = 0$. The energy conservation equation $n^\nu T^\mu_{\nu;\mu} = 0$ for first-order density perturbations and on superhorizon scales give

$$\delta\dot{\rho} = -3H(\delta\rho + \delta p) - 3\dot{\psi}(\rho + p).$$

We write $\delta p = \delta p_{\text{nad}} + c_s^2 \delta\rho$, where δp_{nad} is the non-adiabatic component of the pressure perturbation and $c_s^2 = \delta p_{\text{ad}}/\delta\rho$ is the adiabatic one. In the uniform-density gauge $\psi = \zeta$ and $\delta\rho = 0$ and therefore $\delta p_{\text{ad}} = 0$. The energy conservation equation implies

$$\dot{\zeta} = -\frac{H}{p + \rho} \delta p_{\text{nad}}.$$

If perturbations are adiabatic, the curvature on uniform-density gauge is constant on superhorizon scales. The same holds for the comoving curvature $\mathcal{R}$ as the latter and ζ are equal on superhorizon scales, see section 7.

References

- [1] L. F. Abbott, E. Farhi and M. B. Wise, Phys. Lett. B **117**, 29 (1982).
- [2] V. Acquaviva, N. Bartolo, S. Matarrese and A. Riotto, arXiv:astro-ph/0209156.
- [3] A. Albrecht and P. J. Steinhardt, Phys. Rev. Lett **48**, 1220 (1982).
- [4]
- [4] L. Amendola, C. Gordon, D. Wands and M. Sasaki, Phys. Rev. Lett. **88**, 211302 (2002).
- [5] C. Baccigalupi, A. Balbi, S. Matarrese, F. Perrotta and N. Vittorio, Phys. Rev. D **65**, 063520 (2002).
- [6] For a review, see, for instance, N. A. Bahcall, J. P. Ostriker, S. Perlmutter and P. J. Steinhardt, Science **284**, 1481 (1999).
- [7] J.M. Bardeen, Phys. Rev. D**22**, 1882 (1980); J.M. Bardeen, P. J. Steinhardt and M. S. Turner, Phys. Rev. D**28**, 679 (1983).
- [8] J. D. Barrow and A. R. Liddle, Phys. Rev. D **47**, R5219 (1993).
- [9] N. Bartolo, S. Matarrese and A. Riotto, Phys. Rev. D **64**, 083514 (2001).
- [10] N. Bartolo, S. Matarrese and A. Riotto, Phys. Rev. D **64**, 123504 (2001).
- [11] N. Bartolo, S. Matarrese and A. Riotto, Phys. Rev. D **65**, 103505 (2002).
- [12] N. Bartolo, S. Matarrese, A. Riotto and D. Wands, Phys. Rev. D **66**, 043520 (2002).
- [13] P. Binetruy and G. R. Dvali, Phys. Lett. B **388**, 241 (1996).
- [14] J.R. Bond et *al.*, astro-ph/0210007.
- [15] See <http://www.physics.ucsb.edu/~boomerang>.
- [16] E. J. Copeland, A. R. Liddle, D. H. Lyth, E. D. Stewart, and D. Wands, Phys. Rev. D **49**, 6410 (1994).
- [17] D. J. Chung, E. W. Kolb, A. Riotto and I. I. Tkachev, Phys. Rev. D **62**, 043508 (2000).
- [18] See <http://astro.uchicago.edu/dasi>.
- [19] M. Dine and A. Riotto, Phys. Rev. Lett. **79**, 2632 (1997).
- [20] S. Dodelson, W. H. Kinney and E. W. Kolb, Phys. Rev. D **56** 3207 (1997).
- [21] A. D. Dolgov, Phys. Rept. **222**, 309 (1992).
- [22] G. R. Dvali and A. Riotto, Phys. Lett. B **417**, 20 (1998).
- [23] J. R. Espinosa, A. Riotto and G. G. Ross, Nucl. Phys. B **531**, 461 (1998).

- [24] K. Freese, J. Frieman and A. Olinto, Phys. Rev. Lett. **65**, 3233 (1990).
- [25] W. Friedmann, plenary talk given at *COSMO02*, Chicago, Illinois, USA, September 18-21, 2002.
- [26] A. H. Guth, Phys. Rev. D **23**, 347 (1981).
- [27] A. H. Guth and S.-Y. Pi, Phys. Rev. Lett. **49** 1110 (1982).
- [28] E. Halyo, Phys. Lett. B **387**, 43 (1996).
- [29] S. W. Hawking, Phys. Lett. **B115** 295 (1982) .
- [30] W. Hu, this series of lectures.
- [31] W. H. Kinney and A. Riotto, Astropart. Phys. **10**, 387 (1999), hep-ph/9704388.
- [32] W. H. Kinney and A. Riotto, Phys. Lett. **435B**, 272 (1998).
- [33] W. H. Kinney, A. Melchiorri and A. Riotto, Phys. Rev. D **63**, 023505 (2001).
- [34] S. F. King and A. Riotto, Phys. Lett. B **442**, 68 (1998).
- [35] L. Kofman, A. D. Linde and A. A. Starobinsky, Phys. Rev. Lett. **73**, 3195 (1994).
- [36] E. W. Kolb and M. S. Turner, “The Early universe,” *Redwood City, USA: Addison-Wesley (1990) 547 p. (Frontiers in physics, 69)*.
- [37] E. W. Kolb, arXiv:hep-ph/9910311.
- [38] A.R. Liddle and D. H. Lyth, 1993, Phys. Rept. **231**, 1 (1993).
- [39] A.R. Liddle and D.H. Lyth, *COSMOLOGICAL INFLATION AND LARGE-SCALE STRUCTURE*, Cambridge Univ. Pr. (2000).
- [40] . E. Lidsey, A. R. Liddle, E. W. Kolb, E. J. Copeland, T. Barreiro and M. Abney, Rev. Mod. Phys. **69**, 373 (1997).
- [41] A. D. Linde, *Particle Physics and Inflationary Cosmology*, Harwood Academic, Switzerland (1990).
- [42] A. D. Linde, Phys. Lett. **B108** 389, 1982.
- [43] A. D. Linde, Phys. Lett. **B129**, 177 (1983).
- [44] A. Linde, Phys. Lett. **B259**, 38 (1991).
- [45] A. Linde, Phys. Rev. D **49**, 748 (1994).
- [46] A. D. Linde and A. Riotto, Phys. Rev. D **56**, 1841 (1997).
- [47] D. H. Lyth and A. Riotto, Phys. Lett. B **412**, 28 (1997).

- [48] D. H. Lyth and A. Riotto, Phys. Rept. **314**, 1 (1999).
- [49] J. Mather et al., *Astrophys. J.*, in press (1993).
- [50] See <http://map.gsfc.nasa.gov>.
- [51] See <http://cosmology.berkeley.edu/group/cmb>.
- [52] V. F. Mukhanov, Sov. Phys. JETP **67** (1988) 1297 [Zh. Eksp. Teor. Fiz. **94N7** (1988 ZETFA,94,1-11.1988) 1].
- [53] V.F. Mukhanov, H.A. Feldman and R.H. Brandenberger, Phys. Rept. **215**, 203 (1992).
- [54] For a more complete pedagogical discussion of structure formation see e.g. P.J.E. Peebles, *The Large-scale Structure of the universe* (Princeton Univ. Press, Princeton, 1980)
- [55] P.J.E. Peebles, D.N. Schramm, E. Turner, and R. Kron, *Nature* **352**, 769 (1991).
- [56] See <http://astro.estec.esa.nl/SA-general/Projects/Planck>.
- [57] A. Riotto, Nucl. Phys. B **515**, 413 (1998).
- [58] A. Riotto, arXiv:hep-ph/9710329.
- [59] For a review, see A. Riotto, hep-ph/9807454, lectures delivered at the *ICTP Summer School in High-Energy Physics and Cosmology*, Miramare, Trieste, Italy, 29 Jun - 17 Jul 1998.
- [60] A. Riotto and M. Trodden, Ann. Rev. Nucl. Part. Sci. **49**, 35 (1999).
- [61] R.K. Sachs and A.M. Wolfe, *Astrophys. J.* **147**, 73 (1967).
- [62] M. Sasaki, Prog. Theor. Phys. **76**, 1036 (1986).
- [63] A. A. Starobinsky, Phys. Lett. **B117** 175 (1982) .
- [64] J.M. Stewart, Class. Quant. Grav. **7**, 1169 (1990).
- [65] See <http://www.hep.upenn.edu/~max>.
- [66] D. Wands, K. A. Malik, D. H. Lyth and A. R. Liddle, Phys. Rev. D **62**, 043527 (2000).




Figure 1: The black body spectrum of the cosmic background radiation




Figure 2: The horizon scale (green line) and a physical scale λ (red line) as function of the scale factor a . From Ref. [37].




Figure 3: The CMBR anisotropy as function of ℓ . From Ref. [65].




Figure 4: The behaviour of a generic scale λ and the horizon scale H^{-1} in the standard inflationary model. From Ref. [37].




Figure 5: Large field models of inflation. From Ref. [37].




Figure 6: Small field models of inflation. From Ref. [37].




Figure 7: Hybrid field models of inflation. From Ref. [37].




Figure 8: The horizon scale (green line) and a physical scale λ (red line) as function of the scale factor a . From Ref. [37].




Figure 9: In the reference unperturbed universe, constant-time surfaces have constant spatial curvature (zero for a flat FRW model). In the actual perturbed universe, constant-time surfaces have spatially varying spatial curvature. From Ref. [37].